

Pair of Straight Lines

EXERCISE 4.1 [PAGES 119 - 120]

Exercise 4.1 | Q 1.1 | Page 119

Find the combined equation of the following pair of line:

$$2x + y = 0$$
 and $3x - y = 0$

Solution: The combined equation of the lines 2x + y = 0 and 3x - y = 0 is

$$(2x + y)(3x - y) = 0$$

 $\therefore 6x^2 - 2xy + 3xy - y^2 =$
 $\therefore 6x^2 + xy - y^2 = 0$

Exercise 4.1 | Q 1.2 | Page 119

Find the combined equation of the following pair of line:

x + 2y - 1 = 0 and x - 3y + 2 = 0

Solution: The combined equation of the lines x + 2y - 1 = 0 and x - 3y + 2 = 0 is

$$(x + 2y - 1)(x - 3y + 2) = 0$$

 $\therefore x^2 - 3xy + 2x + 2xy - 6y^2 + 4y - x + 3y - 2 = 0$

0

 $\therefore x^2 - xy - 6y^2 + x + 7y - 2 = 0$

Exercise 4.1 | Q 1.3 | Page 119

Find the combined equation of the following pair of line:

passing through (2, 3) and parallel to the coordinate axes.

Solution: Equations of the coordinate axes are x = 0 and y = 0

: The equations of the lines passing through (2, 3) and parallel to the coordinate axes are x = 2 and y = 3.

i.e. x - 2 = 0 and y - 3 = 0

 \therefore their combined equation is

$$(x - 2)(y - 3) = 0$$

 $\therefore xy - 3x - 2y + 6 = 0$

Exercise 4.1 | Q 1.4 | Page 119



Find the combined equation of the following pair of line:

passing through (2, 3) and perpendicular to the lines 3x + 2y - 1 = 0 and x - 3y + 2 = 0

Solution: Let L_1 and L_2 be the lines passing through the point (2, 3) and perpendicular to the lines 3x + 2y - 1 = 0 and x - 3y + 2 = 0 respectively.

Slopes of the lines 3x + 2y - 1 = 0 and x - 3y + 2 = 0 are $\frac{-3}{2}$ and $\frac{-1}{-3} = \frac{1}{3}$ respectively.

 \therefore slopes of the lines L₁ and L₂ pass through the point (2, 3), their equations are

y - 3 = 2/3(x - 2) and y - 3 = -3(x - 2) ∴ 3y - 9 = 2x - 4 and y - 3 = - 3x + 6 ∴ 2x - 3y + 5 = 0 and 3x + y - 9 = 0 their combined equation is (2x - 3y + 5)(3x + y - 9) = 0

 $\therefore 6x^2 + 2xy - 18x - 9xy - 3y^2 + 27y + 15x + 5y - 45 = 0$

 $\therefore 6x^2 - 7xy - 3y^2 - 3x + 32y - 45 = 0$

Exercise 4.1 | Q 1.5 | Page 119

Find the combined equation of the following pair of line:

passing through (-1, 2), one is parallel to x + 3y - 1 = 0 and other is perpendicular to 2x - 3y - 1 = 0

Solution: Let L_1 be the line passing through the point (-1, 2) and parallel to the line x + 3y - 1 = 0 whose slope is -1/3.

- \therefore slope of the line L₁ is 1/3
- \therefore equation of the line L_1 is
- y 2 = -1/3 (x + 1)
- ∴ 3y 6 = x 1
- $\therefore x + 3y 5 = 0$

Let L_1 be the line passing through (-1, 2) and perpendicular to the line 2x - 3y - 1 = 0 whose slope is

$$\frac{-2}{-3} = \frac{2}{3}$$

- \therefore slope of the line $L_2\ is-1/3$
- \therefore equation of the line L₂ is



y - 2 = - 1/3 (x + 1) ∴ 2y - 4 = - 3x - 3 ∴ 3x + 2y - 1 = 0 Hence, the equations of the required lines are x + 3y - 5 = 0 and 3x + 2y - 1 = 0 ∴ their combined equation is (x + 3y - 5)(3x + 2y - 1) = 0 ∴ 3x² + 2xy - x + 9xy + 6y² - 3y - 15x - 10y + 5 = 0 ∴ 3x² + 11xy + 6y² - 16x - 13y + 5 = 0

Exercise 4.1 | Q 2.1 | Page 119

Find the separate equation of the line represented by the following equation:

 $3y^2 + 7xy = 0$ Solution: $3y^2 + 7xy = 0$ ∴ y(3y + 7x) = 0∴ the separate equations of the lines are y = 0 and 7x + 3y = 0

Exercise 4.1 | Q 2.2 | Page 119

Find the separate equation of the line represented by the following equation:

$$5y^2 + 9y^2 = 0$$

Solution:

$$5y^{2} + 9y^{2} = 0$$

$$\therefore \left(\sqrt{5x}\right)^{2} - \left(\sqrt{3y}\right)^{2} = 0$$

$$\therefore \left(\sqrt{5x} + 3y\right)\left(\sqrt{5x} - 3y\right) = 0$$

the separate equations of the lines are $\left(\sqrt{5\mathrm{x}}+3\mathrm{y}
ight)=0$ and $\left(\sqrt{5\mathrm{x}}-3\mathrm{y}
ight)=0$

Exercise 4.1 | Q 2.3 | Page 119

Find the separate equation of the line represented by the following equation: $x^2 - 4xy = 0$



Solution: $x^2 - 4xy = 0$

$$\therefore x (x - 4y) = 0$$

: the separate equations of the lines are x = 0 and x - 4y = 0

Exercise 4.1 | Q 2.4 | Page 119

Find the separate equations of the lines represented by the equation $3x^2-10xy-8y^2=0$

Solution:

Given pairs of lines $3x^2 - 10xy - 8y^2 = 0$

$$3x^{2} - 12xy + 2xy - 8y^{2} = 0$$

 $3x(x - 4y) + 2y(x - 4y) = 0$

$$(\mathbf{x} - 4\mathbf{y})(3\mathbf{x} + 2\mathbf{y}) = 0$$

Separated equations

$$3x + 2y = 0$$
 and $x - 4y = 0$

Exercise 4.1 | Q 2.5 | Page 119

Find the separate equation of the line represented by the following equation:

$$3x^2 - 2\sqrt{3}xy - 3y^2 = 0$$

Solution:

$$3x^{2} - 2\sqrt{3}xy - 3y^{2} = 0$$

$$\therefore 3x^{2} - 3\sqrt{3}xy + \sqrt{3}xy - 3y^{2} = 0$$

$$\therefore 3x\left(x - \sqrt{3}y\right) + \sqrt{3}y\left(x - \sqrt{3}y\right) = 0$$

$$\therefore \left(x - \sqrt{3}y\right)\left(3x + \sqrt{3}y\right) = 0$$

The separate equations of the lines are

$$\mathrm{x}-\sqrt{3}\mathrm{y}=0$$
 and $3\mathrm{x}+\sqrt{3}\mathrm{y}=0$

Exercise 4.1 | Q 2.6 | Page 119

Find the separate equation of the line represented by the following equation:



$$x^{2} + 2(\operatorname{cosec} \alpha)xy + y^{2} = 0$$

Solution: $x^{2} + 2(\operatorname{cosec} \alpha)xy + y^{2} = 0$
i.e. $y^{2} + 2(\operatorname{cosec} \alpha)xy + x^{2} = 0$
Dividing by x^{2} , we get,
 $\left(\frac{y}{x}\right)^{2} + 2\operatorname{cosec} \alpha$. $\left(\frac{y}{x}\right) + 1 = 0$
 $\therefore \frac{y}{x} = \frac{-2\operatorname{cosec} \alpha \pm \sqrt{4\operatorname{cosec}^{2}\alpha - 4 \times 1 \times 1}}{2 \times 1}$
 $= \frac{-2\operatorname{cosec} \alpha \pm 2\sqrt{\operatorname{cosec}^{2}\alpha - 1}}{2}$
 $= -\operatorname{cosec} \alpha \pm \cot \alpha$
 $\therefore \frac{y}{x} = (\cot \alpha - \operatorname{cosec} \alpha) \text{ and}$
 $\frac{y}{x} = -(\operatorname{cosec} \alpha + \cot \alpha)$

The separate equations of the lines are

 $(\csc \alpha - \cot \alpha) x + y = 0$ and $(\csc \alpha - \cot \alpha) x + y = 0$

Exercise 4.1 | Q 2.7 | Page 119

Find the separate equation of the line represented by the following equation:

 x^2 + 2xy tan α - y^2 = 0

Solution:

$$x^2$$
 + 2xy tan α - y^2 = 0

Dividing by y²

$$\left(rac{\mathrm{x}}{\mathrm{y}}
ight)^2 + 2\left(rac{\mathrm{x}}{\mathrm{y}}
ight) \mathrm{tan}lpha - 1 = 0$$



$$\therefore \frac{\mathbf{x}}{\mathbf{y}} = \frac{-2\tan\alpha \pm \sqrt{4\tan^2\alpha - 4 \times 1 \times 1}}{2 \times 1}$$
$$= \frac{-2\tan\alpha \pm 2\sqrt{\tan^2\alpha - 1}}{2}$$
$$= -\tan\alpha \pm \sec\alpha$$
$$\left(\frac{\mathbf{x}}{\mathbf{y}}\right) = (\sec\alpha - \tan\alpha) \text{ and}$$
$$\left(\frac{\mathbf{x}}{\mathbf{y}}\right) = -(\tan\alpha + \sec\alpha)$$

The separate equations of the lines are

 $(\sec \alpha - \tan \alpha) x + y = 0$ and $(\sec \alpha + \tan \alpha)x - y = 0$

Exercise 4.1 | Q 3.1 | Page 119

Find the combined equation of the pair of a line passing through the origin and perpendicular to the line represented by following equation:

$$5x^2 - 8xy + 3y^2 = 0$$

Solution: Comparing the equation $5x^2 - 8xy + 3y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = 5, 2h = - 8, b= 3

Let m_1 and m_2 be the slopes of the lines represented by $5x^2 - 8xy + 3y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{8}{3}$$
 and $m_1m_2 = \frac{a}{b} = \frac{5}{3}$ (1)

Now required lines are perpendicular to these lines

$$\therefore$$
 their slopes are $\displaystyle\frac{-1}{m_1}$ and $\displaystyle\frac{-1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$\mathsf{y} = rac{-1}{\mathbf{m}_1} \mathbf{x} ext{ and } \mathsf{y} = rac{-1}{\mathbf{m}_2} \mathbf{x}$$



i.e. $m_1y = -x$ and $m_2y = -x$

i.e. $x + m_1 y = 0$ and $x + m_2 y = 0$

 \therefore their combined equation is

$$(x + m_1 y)(x + m_2 y) = 0$$

$$\therefore x^{2} + (m_{1} + m_{2})xy + m_{1}m_{2}y^{2} = 0$$

$$\therefore x^{2} + \frac{8}{3}xy + \frac{5}{3}y^{2} = 0 \quad ... [By (1)]$$

$$\therefore 3x^2 + 8xy + 5y^2 = 0$$

Exercise 4.1 | Q 3.2 | Page 119

Find the combined equation of the pair of a line passing through the origin and perpendicular to the line represented by the following equation:

$$5x^2 + 2xy - 3y^2 = 0$$

Solution: Comparing the equation $5x^2 + 2xy - 3y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,

Let m_1 and m_2 be the slopes of the lines represented by $5x^2 + 2xy - 3y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{-2}{-3} = \frac{2}{3} \text{ and } m_1m_2 = \frac{a}{b} = \frac{5}{-3}$$
(1)

Now required lines are perpendicular to these lines

$$\therefore$$
 their slopes are $\displaystyle\frac{-1}{m_1}$ and $\displaystyle\frac{-1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$\mathsf{y} = \frac{-1}{\mathbf{m}_1} \mathbf{x} \text{ and } \mathsf{y} = \frac{-1}{\mathbf{m}_2} \mathbf{x}$$



i.e.
$$m_1y = -x$$
 and $m_2y = -x$

i.e.
$$x + m_1 y = 0$$
 and $x + m_2 y = 0$

 \therefore their combined equation is

$$(x + m_1 y)(x + m_2 y) = 0$$

$$\therefore x^2 + (m_1 + m_2)xy + m_1m_2y^2 = 0$$

∴
$$\mathbf{x}^2 + \frac{2}{3}\mathbf{x}\mathbf{y} - \frac{5}{3}\mathbf{y}^2 = 0$$
 ...[By (1)]
∴ $3\mathbf{x}^2 + 2\mathbf{x}\mathbf{y} - 5\mathbf{y}^2 = 0$

Exercise 4.1 | Q 3.3 | Page 119

Find the combined equation of the pair of a line passing through the origin and perpendicular to the line represented by the following equation:

$$xy + y^2 = 0$$

Solution: Comparing the equation $xy + y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,

a = 0, 2h = 1, b= 1

Let m_1 and m_2 be the slopes of the lines represented by $xy + y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{-1}{1} = -1 \text{ and } m_1m_2 = \frac{a}{b} = \frac{0}{1} = 0 \quad(1)$$

Now required lines are perpendicular to these lines

$$\therefore$$
 their slopes are $\displaystyle\frac{-1}{m_1}$ and $\displaystyle\frac{-1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$y = \frac{-1}{m_1}x$$
 and $y = \frac{-1}{m_2}x$

i.e. $m_1y = -x$ and $m_2y = -x$ i.e. $x + m_1y = 0$ and $x + m_2y = 0$



 \therefore their combined equation is

 $(x + m_1 y)(x + m_2 y) = 0$

 $\therefore x^2 + (m_1 + m_2)xy + m_1m_2y^2 = 0$

$$\therefore x^2 - xy + 0. y^2 = 0$$
 ...[By (1)]

 $\therefore x^2 - xy = 0$

[Note: : Answer in the textbook is incorrect.]

Exercise 4.1 | Q 3.4 | Page 119

Find the combined equation of the pair of a line passing through the origin and perpendicular to the line represented by the following equation:

 $3x^2 - 4xy = 0$

Solution: Consider $3x^2 - 4xy = 0$

$$\therefore x(3x - 4y) = 0$$

 \therefore separate equations of the lines are x = 0 and 3x - 4y = 0

Let m₁ and m₂ be the slopes of these lines.

Then m₁ does not exist and m₂ = $\frac{3}{4}$

Now, required lines are perpendicular to these lines.

:. their slopes are
$$-\frac{1}{m_1}$$
 and $-\frac{1}{m_2}$
Since m_1 does not exist, $-\frac{1}{m_1} = 0$
Also, $m_2 = \frac{3}{4}, -\frac{1}{m_2} = -\frac{4}{3}$

Since these lines are passing through the origin, their

separate equations are y = 0 and y = $-\frac{4}{3}x$, i.e. 4x + 3y = 0

 \therefore their combined equation is



y(4x + 3y) = 0 $\therefore 4xy + 3y^2 = 0$

Exercise 4.1 | Q 4.1 | Page 119

Find k, if the sum of the slopes of the lines represented by $x^2 + kxy - 3y^2 = 0$ is twice their product.

Solution: Comparing the equation $x^2 + kxy - 3y^2 = 0$ with $ax^2 + 2hxy - by^2 = 0$, we get, a = 1, 2h = k, b = -3.

Let m_1 and m_2 be the slopes of the lines represented by $x^2 + kxy - 3y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = -\frac{k}{-3} = \frac{k}{3}$$

and $m_1m_2 = \frac{a}{b} = \frac{1}{-3} = -\frac{1}{3}$

Now, $m_1 + m_2 = 2(m_1m_2)$...(given)

$$\therefore \frac{\mathbf{k}}{3} = 2\left(-\frac{1}{3}\right)$$

Exercise 4.1 | Q 4.2 | Page 119

Find k, the slopes of the lines represented by $3x^2 + kxy - y^2 = 0$ differ by 4. **Solution:** Comparing the equation $3x^2 + kxy - y^2 = 0$ with $ax^2 + 2hxy - by^2 = 0$, we get, a = 3, 2h = k, b = -1.

Let m_1 and m_2 be the slopes of the lines represented by $3x^2 + kxy - y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = -\frac{k}{-1} = k$$

and $m_1m_2 = \frac{a}{b} = \frac{3}{-1} = -3$
$$\therefore (m_1 - m_2)^2 = (m_1 + m_2)^2 - 4m_1m_2$$

$$= k^2 - 4 (-3)$$

$$= k^2 + 12 \qquad \dots (1)$$



But $|m_1 - m_2| = 4$ $\therefore (m_1 - m_2)^2 = 16 \dots (2)$ \therefore from (1) and (2), $k^2 + 12 = 16$ $\therefore k^2 = 4$ $\therefore k = \pm 2$

Exercise 4.1 | Q 4.3 | Page 119

Find k, the slope of one of the lines given by $kx^2 + 4xy - y^2 = 0$ exceeds the slope of the other by 8.

Solution: Comparing the equation $kx^2 + 4xy - y^2 = 0$ with $ax^2 + 2hxy - by^2 = 0$, we get, a = k, 2h = 4, b = -1.

Let m_1 and m_2 be the slopes of the lines represented by $kx^2 + 4xy - y^2 = 0$

$$\begin{array}{l} \therefore \ m_1 + m_2 = \displaystyle \frac{-2h}{b} = \displaystyle -\frac{4}{-1} = 4 \\ \text{and} \ m_1 m_2 = \displaystyle \frac{a}{b} = \displaystyle \frac{k}{-1} = \displaystyle -k \\ \text{We are given that} \ m_2 = m_1 + 8 \\ \therefore \ m_1 + m_1 + 8 = 4 \\ \therefore \ 2m_1 = -4 \quad \therefore \ m_1 = -2 \quad \dots(1) \\ \text{Also,} \ m_1(m_1 + 8) = -k \\ (-2)(-2 + 8) = -k \quad \dots[By\ (1)] \\ \therefore \ (-2)(6) = -k \\ \therefore -12 = -k \\ \therefore \ k = 12 \end{array}$$

Exercise 4.1 | Q 5.1 | Page 120

Find the condition that the line 4x + 5y = 0 coincides with one of the lines given by $ax^2 + bx^2 + bx^$

 $2hxy + by^2 = 0$

Solution: The auxiliary equation of the lines represented by $ax^2 + 2hxy + by^2 = 0$ is $bm^2 + 2hm + a = 0$

Given that 4x + 5y = 0 is one of the lines represented by $ax^2 + 2hxy + by^2 = 0$



The slope of the line 4x + 5y = 0 is $-\frac{4}{5}$

$$\therefore m = -\frac{4}{5} \text{ is a root of the auxiliary equation } bm^{2} + 2hm + a = 0$$

$$\therefore b\left(-\frac{4}{5}\right)^{2} + 2h\left(-\frac{4}{5}\right) + a = 0$$

$$\therefore \frac{16b}{25} - \frac{8h}{5} + a = 0$$

$$\therefore 16b - 40h + 25a = 0$$

$$\therefore 25a + 16b = 40h$$

This is the required condition.

Exercise 4.1 | Q 5.2 | Page 120

Find the condition that the line 3x + y = 0 may be perpendicular to one of the lines given by $ax^2 + 2hxy + by^2 = 0$

Solution: The auxiliary equation of the lines represented by $ax^2 + 2hxy + by^2 = 0$ is $bm^2 + 2hm + a = 0$

Since one line is perpendicular to the line 3x + y = 0 whose slope is $-\frac{3}{1} = -3$

 $\therefore \text{ slope of that line } = m = \frac{1}{3}$ $\therefore m = \frac{1}{3} \text{ is the root of the auxiliary equation } bm^2 + 2hm + a = 0.$ $\therefore b\left(\frac{1}{3}\right)^2 + 2h\left(\frac{1}{3}\right) + a = 0$ $\therefore \frac{b}{9} + \frac{2h}{3} + a = 0$ $\therefore b + 6h + 9a = 0$ $\therefore 9a + b + 6h = 0$

This is the required condition.



Exercise 4.1 | Q 6 | Page 120

If one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is perpendicular to px + qy = 0, show that $ap^2 + 2hpq + bq^2 = 0$.

Solution: To prove: $ap^2 + 2hpq + bq^2 = 0$.

Let the slope of the pair of straight lines $ax^2 + 2hxy + by^2 = 0$ be m₁ and m₂

Then,
$$m_1 + m_2 = \frac{-2h}{b}$$
 and $m_1m_2 = \frac{a}{b}$

Slope of the line px + qy = 0 is $\frac{-p}{q}$

But one of the lines of $ax^2 + 2hxy + by^2 = 0$ is perpendicular to px + qy = 0

$$\Rightarrow \mathbf{m}_{1} = \frac{\mathbf{q}}{\mathbf{p}}$$
Now, $\mathbf{m}_{1} + \mathbf{m}_{2} = \frac{-2\mathbf{h}}{\mathbf{b}}$ and $\mathbf{m}_{1}\mathbf{m}_{2} = \frac{\mathbf{a}}{\mathbf{b}}$

$$\Rightarrow \frac{\mathbf{q}}{\mathbf{p}} + \mathbf{m}_{2} = \frac{-2\mathbf{h}}{\mathbf{b}}$$
 and $\left(\frac{\mathbf{q}}{\mathbf{p}}\right)\mathbf{m}_{2} = \frac{\mathbf{a}}{\mathbf{b}}$

$$\Rightarrow \frac{\mathbf{q}}{\mathbf{p}} + \mathbf{m}_{2} = \frac{-2\mathbf{h}}{\mathbf{b}}$$
 and $\mathbf{m}_{2} = \frac{\mathbf{a}\mathbf{p}}{\mathbf{b}\mathbf{q}}$

$$\Rightarrow \frac{\mathbf{q}}{\mathbf{p}} + \frac{\mathbf{a}\mathbf{p}}{\mathbf{b}\mathbf{q}} = \frac{-2\mathbf{h}}{\mathbf{b}}$$

$$\Rightarrow \frac{\mathbf{b}\mathbf{q}^{2} + \mathbf{a}\mathbf{p}^{2}}{\mathbf{p}\mathbf{q}} = -2\mathbf{h}$$

$$\Rightarrow \mathbf{b}\mathbf{q}^{2} + \mathbf{a}\mathbf{p}^{2} = -2\mathbf{h}\mathbf{p}\mathbf{q}$$

$$\Rightarrow \mathbf{a}\mathbf{p}^{2} + 2\mathbf{h}\mathbf{p}\mathbf{q} + \mathbf{b}\mathbf{q}^{2} = 0$$

Exercise 4.1 | Q 7 | Page 120

Find the combined equation of the pair of lines through the origin and making an equilateral triangle with the line y = 3.

Solution: Let OA and OB be the lines through the origin making an angle of 60° with the line y = 3.

 \therefore OA and OB make an angle of 60° and 120° with the positive direction of the X-axis.



∴ slope of OA = tan 60° = $\sqrt{3}$

 \therefore equation of the line OA is



Slope of OB = $\tan 120^\circ = \tan (180^\circ - 60^\circ)$

= - tan 60° = - $\sqrt{3}$

∴ equation of the line OB is

y =
$$-\sqrt{3}x$$
, i.e. $\sqrt{3}x + y = 0$

 \therefore required joint equation of the lines is

$$\left(\sqrt{3}x - y\right)\left(\sqrt{3}x + y\right) = 0$$

i.e. $3x^2 - y^2 = 0$

Exercise 4.1 | Q 8 | Page 120

If the slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is four times the other, show that $16h^2 = 25ab$.

Solution: Let m_1 and m_2 be the slopes of the lines given by $ax^2 + 2hxy + by^2 = 0$



$$\therefore m_1 + m_2 = -\frac{2h}{b}$$

and $m_1m_2 = \frac{a}{b}$

We are given that $m_2 = 4m_1$

$$\therefore m_{1} + 4m_{1} = -\frac{2h}{b}$$

$$\therefore 5m_{1} = \frac{-2h}{b}$$

$$\therefore m_{1} = -\frac{2h}{5b} \quad \dots (1)$$
Also, $m_{1}(4m_{1}) = \frac{a}{b}$

$$\therefore 4m_{1}^{2} = \frac{a}{b}$$

$$\therefore m_{1}^{2} = \frac{a}{4b}$$

$$\therefore \left(\frac{-2h}{5b}\right)^{2} = \frac{a}{4b} \quad \dots [By(1)]$$

$$\therefore \frac{4h^{2}}{25b^{2}} = \frac{a}{4b}$$

$$\therefore \frac{4h^{2}}{25b} = \frac{a}{4}, \text{ as } b \neq 0$$

$$\therefore 16h^{2} = 25ab$$

This is the required condition.

Exercise 4.1 | Q 9 | Page 120



If one of the lines given by $ax^2 + 2hxy + by^2 = 0$ bisect an angle between the coordinate axes, then show that $(a + b)^2 = 4h^2$.

Solution: The auxiliary equation of the lines given by $ax^2 + 2hxy + by^2 = 0$ is $bm^2 + 2hm + a = 0$.

Since one of the lines bisects an angle between the coordinate axes, that line makes an angle of 45° or 135° with the positive direction of X-axis.

 \therefore the slope of that line = tan 45° or tan 135°

∴ m = tan 45° = 1

or m = tan 135° = tan ($180^{\circ} - 45^{\circ}$)

= - tan 45° = - 1

 \therefore m = ± 1 are the roots of the auxiliary equation bm² + 2hm + a = 0.

$$b(\pm 1)^2 + 2h(\pm 1) + a = 0$$

 \therefore b ± 2h + a = 0

 \therefore a + b = ± 2h

$$(a + b)^2 = 4h^2$$

This is the required condition.

EXERCISE 4.2 [PAGE 124]

Exercise 4.2 | Q 1 | Page 124

. Show that the lines represented by $3x^2 - 4xy - 3y^2 = 0$ are perpendicular to each other.

Solution: Comparing the equation $3x^2 - 4xy - 3y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = 3, 2h = -4, b = -3.

Since a + b = 3 + (-3) = 0, the lines represented by $3x^2 - 4xy - 3y^2 = 0$ are perpendicular to each other.

Exercise 4.2 | Q 2 | Page 124

Show that the lines represented by $x^2 + 6xy + 9y^2 = 0$ are coincident.

Solution: Comparing the equation $x^2 + 6xy + 9y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = 1, 2h = 6 i.e. h = 3 and b = 9.

Since $h^2 - ab = (3)^2 - 1(9)$ = 9 - 9 = 0, the lines represented by $x^2 + 6xy + 9y^2 = 0$ are coincident.

Exercise 4.2 | Q 3 | Page 124



Find the value of k if lines represented by $kx^2 + 4xy - 4y^2 = 0$ are perpendicular to each other.

Solution: Comparing the equation $kx^2 + 4xy - 4y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = k, 2h = 4 and b = -4.

Since lines represented by $kx^2 + 4xy - 4y^2 = 0$ are perpendicular to each other,

a + b = 0

∴ k - 4 = 0

∴ k = 4

Exercise 4.2 | Q 4.1 | Page 124

Find the measure of the acute angle between the line represented by: $3x^2 - 4\sqrt{3}xy + 3y^2 = 0$

Solution:

Comparing the equation

$$3x^2 - 4\sqrt{3}xy + 3y^2 = 0$$
 with
ax² + 2hxy + by² = 0, we get,
a = 3, 2h = -4\sqrt{3} i.e. h = $-2\sqrt{3}$ and b = 3

Let θ be the acute angle between the lines.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$= \left| \frac{2\sqrt{\left(-2\sqrt{3}\right)^2 - 3(3)}}{3 + 3} \right|$$



$$= \left| \frac{2\sqrt{12 - 9}}{6} \right|$$
$$= \left| \frac{2\sqrt{3}}{6} \right|$$
$$\therefore \tan \theta = \frac{1}{\sqrt{3}} = \tan 30^{\circ}$$

$$\therefore \theta = 30^{\circ}$$

Exercise 4.2 | Q 4.2 | Page 124

Find the measure of the acute angle between the line represented by:

$$4x^2 + 5xy + y^2 = 0$$

Solution: Comparing the equation

$$4x^2 + 5xy + y^2 = 0$$
 with

 $ax^{2} + 2hxy + by^{2} = 0$, we get,

a = 4, 2h = 5 i.e. h = 5/2 and b = 1

Let θ be the acute angle between the lines.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$= \left| \frac{2\sqrt{\left(\frac{5}{2}\right)^2 - 4(1)}}{4 + 1} \right|$$



$$= \left| \frac{2\sqrt{\left(\frac{25}{4}\right) - 4}}{5} \right|$$
$$= \left| \frac{2 \times \frac{3}{2}}{5} \right|$$
$$\therefore \tan \theta = \frac{3}{5}$$
$$\therefore \theta = \tan^{-1} \left(\frac{3}{5} \right)$$

Exercise 4.2 | Q 4.3 | Page 124

Find the measure of the acute angle between the line represented by:

$$2x^2 + 7xy + 3y^2 = 0$$

Solution: Comparing the equation

$$2x^2 + 7xy + 3y^2 = 0$$
 with

 $ax^{2} + 2hxy + by^{2} = 0$, we get,

$$a = 2$$
, $2h = 7$ i.e. $h = 7/2$ and $b = 3$

Let θ be the acute angle between the lines.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$= \left| \frac{2\sqrt{\left(\frac{7}{2}\right)^2 - 2(3)}}{2 + 3} \right|$$
$$= \left| \frac{2\sqrt{\left(\frac{49}{4}\right) - 6}}{5} \right|$$





 $\tan \theta = 1$

 $\therefore \theta = \tan 1 = 45^{\circ}$

∴ θ = 45°

Exercise 4.2 | Q 4.4 | Page 124

Find the measure of the acute angle between the line represented by:

```
(a^{2} + 3b^{2})x^{2} + 8abxy + (b^{2} - 3a^{2})y^{2} = 0
Solution: Comparing the equation
(a^{2} + 3b^{2})x^{2} + 8abxy + (b^{2} - 3a^{2})y^{2} = 0 \text{ with}Ax^{2} + 2Hxy + By^{2} = 0, \text{ we have,}A = a^{2} + 3b^{2}, H = 4ab \text{ and } B = b^{2} - 3a^{2}\therefore H^{2} - AB = 16a^{2}b^{2} - (a^{2} - 3b^{2})(b^{2} - 3a^{2})= 16a^{2}b^{2} + (a^{2} - 3b^{2})(3a^{2} - b^{2})= 16a^{2}b^{2} + 3a^{4} - 10a^{2}b^{2} + 3b^{4}= 3a^{4} + 6a^{2}b^{2} + 3b^{4}= 3(a^{4} + 2a^{2}b^{2} + b^{4})= 3(a^{2} + b^{2})^{2}
```



$$\therefore \sqrt{H^2 - AB} = \sqrt{3}(a^2 + b^2)$$

Also, A + B = (a² - 3b²) + (b² - 3a²)
= - 2(a² + b²)

Let θ be the acute angle between the lines, then

$$\therefore \tan \theta = \left| \frac{2\sqrt{H^2 - AB}}{A + B} \right|$$
$$= \left| \frac{2\sqrt{3}(a^2 + b^2)}{-2(a^2 + b^2)} \right|$$
$$= \sqrt{3} = \tan 60^\circ$$
$$\therefore \theta = 60^\circ$$

Exercise 4.2 | Q 5 | Page 124

Find the combined equation of lines passing through the origin each of which making an angle of 30° with the line 3x + 2y - 11 = 0

Solution: The slope of the line 3x + 2y - 11 = 0 is $m_1 = -3/2$

Let m be the slope of one of the lines making an angle of 30° with the line 3x + 2y - 11 = 0

The angle between the lines having slopes m and m_1 is 30°.

$$\therefore \tan 30^\circ = \left| \frac{\mathrm{m} - \mathrm{m}_1}{1 + \mathrm{m} \cdot \mathrm{m}_1} \right|, \text{ where } \tan 30^\circ = \frac{1}{\sqrt{30}}$$
$$\therefore \frac{1}{\sqrt{30}} = \left| \frac{\mathrm{m} - \left(-\frac{3}{2}\right)}{1 + \mathrm{m}\left(-\frac{3}{2}\right)} \right|$$
$$\therefore \frac{1}{\sqrt{30}} = \left| \frac{2\mathrm{m} + 3}{2 - 3\mathrm{m}} \right|$$



On squaring both sides, we get,

$$\frac{1}{3} = \frac{(2m+3)^2}{(2-3m)^2}$$

$$\therefore (2 - 3m)^2 = 3(2m+2)^2$$

$$\therefore 4 - 12m + 9m^2 = 3(4m^2 + 12m + 9)$$

$$\therefore 4 - 12m + 9m^2 = 12m^2 + 36m + 27$$

$$\therefore 3m^2 + 48m + 23 = 0$$

This is the auxiliary equation of the two lines and their joint equation is obtained by putting m = y/x

 \therefore the combined equation of the two lines is

$$3\left(\frac{y}{x}\right)^2 + 48\left(\frac{y}{x}\right) + 23 = 0$$

$$\therefore \frac{3y^2}{x^2} + \frac{48y}{x} + 23 = 0$$

$$\therefore 3y^2 + 48xy + 23x^2 = 0$$

$$\therefore 23x^2 + 48xy + 3y^2 = 0$$

Exercise 4.2 | Q 6 | Page 124

If the angle between the lines represented by $ax^2 + 2hxy + by^2 = 0$ is equal to the angle between the lines $2x^2 - 5xy + 3y^2 = 0$, then show that $100 (h^2 - ab) = (a + b)^2$.

Solution:

The acute angle θ between the lines ax² + 2hxy + by² = 0 is given by

$$\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| \dots (1)$$

Comparing the equation $2x^2 - 5xy + 3y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get. a = 2, 2h = -5, i.e. h = $-\frac{5}{2}$ and b = 3



Let α be the acute angle between the lines $2x^2 - 5xy + 3y^2 = 0$

 $\therefore \tan \alpha = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$ $= \left| \frac{2\sqrt{\left(\frac{5}{2}\right)^2 - 2(3)}}{2 + 3} \right|$ $= \left| \frac{\left(2\frac{\sqrt{25}}{4} - 6\right)}{5} \right|$ $= \left| \frac{2 \times \frac{1}{2}}{5} \right|$ $\therefore \tan \alpha = \frac{1}{2} \qquad (2)$

$$\therefore \tan \alpha = \frac{1}{5}$$
(2)

If
$$\theta = \alpha$$
, then tan $\theta = \tan \alpha$

$$\therefore \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| = \frac{1}{5} \quad \dots [By (1) \text{ and } (2)]$$
$$\therefore \frac{4(h^2 - ab)}{(a + b)^2} = \frac{1}{25}$$

:
$$100 (h^2 - ab) = (a + b)^2$$

This is the required condition.

Exercise 4.2 | Q 7 | Page 124

Find the combined equation of lines passing through the origin and each of which making an angle of 60° with the Y-axis.

Solution:





Let OA and OB be the lines through the origin making an angle of 60° with the Y-axis. Then OA and OB make an angle of 30° and 150° with the positive direction of X-axis.

$$\therefore$$
 slope of OA = tan 30° = $\frac{1}{\sqrt{3}}$

 \therefore equation of the line OA is

$$y = \frac{1}{\sqrt{3}}$$
 x i.e. $x - \sqrt{3}y = 0$

Slope of OB = $\tan 150^\circ = \tan (180^\circ - 30^\circ)$

= - tan 30° =
$$-\frac{1}{\sqrt{3}}$$

 \therefore equation of the line OB is

$$y = -\frac{1}{\sqrt{3}}x$$
 i.e. $x + \sqrt{3}y = 0$

 \therefore required combined equation is

$$\left(x-\sqrt{3}y\right)\left(x+\sqrt{3}y\right)=0$$

i.e. $x^2 - 3y^2 = 0$

EXERCISE 4.3 [PAGES 127 - 128]



Exercise 4.3 | Q 1.1 | Page 127

Find the joint equation of the pair of the line through the point (2, -1) and parallel to the lines represented by $2x^2 + 3xy - 9y^2 = 0$.

Solution: The combined equation of the given lines is

$$2x^{2} + 3xy - 9y^{2} = 0$$

i.e. $2x^{2} + 6xy - 3xy - 9y^{2} = 0$
i.e. $2x(x + 3y) - 3y(x + 3y) = 0$
i.e. $(x + 3y)(2x - 3y) = 0$
∴ their separate equations are
 $x + 3y = 0$ and $2x - 3y = 0$
∴ their slopes are $m_{1} = \frac{-1}{-3}$ and $m_{2} = \frac{-2}{-3} = \frac{2}{3}$
The slopes of the lines parallel to these lines are m_{1} and m_{2} i.e. $-\frac{1}{3}$ and $\frac{2}{3}$.
∴ the equations of the lines with these slopes and through the point (2, -1) are

y + 1 = $-\frac{1}{3}$ (x - 2) and y + 1 = $\frac{1}{3}$ (x - 2) i.e. 3y + 3 = - x + 2 and 3y + 3 = 2x - 4 i.e. x + 3y + 1 = 0 and 2x - 3y - 7 = 0 ∴ the joint equation of these lines is (x + 3y + 1)(2x - 3y - 7) = 0 ∴ 2x² - 3xy - 7x + 6xy - 9y² - 21y + 2x - 3y - 7 = 0 ∴ 2x² + 3xy - 9y² - 5x - 24y - 7 = 0

Exercise 4.3 | Q 1.2 | Page 127

Find the joint equation of the pair of the line through the point (2, -3) and parallel to the lines represented by $x^2 + xy - y^2 = 0$.

Solution: The combined equation of the given lines is

 $x^{2} + xy - y^{2} = 0$ (1) with $ax^{2} + 2hxy + by^{2} = 0$, we get, a = 1, 2h = 1, b = -1



Let m_1 and m_2 be the slopes of the lines represented by (1).

Then
$$m_1 + m_2 = -\frac{2h}{b} = \frac{-1}{-1} = 1$$
 and $m_1m_2 = \frac{a}{b} = \frac{1}{-1} = -1$ (2)

The slopes of the lines parallel to these lines are m_1 and m_2 .

∴ the equations of the lines with these slopes and through the point (2, - 3) are y + 3 = m₁(x - 2) and y + 3 = m₂ (x - 2) i.e. m₁(x - 2) - (y + 3) = 0 and m₂(x - 2) - (y + 3) = 0 ∴ the joint equation of these lines is [m₁ (x - 2) - (y + 3)][m₂(x - 2) - (y + 3)] = 0 ∴ m₁m₂ (x - 2)² - m₁ (x - 2)(y + 3) - m₂(x - 2)(y+3) + (y + 3)² = 0 ∴ m₁m₂ (x - 2)² - (m₁ + m₂)(x - 2)(y + 3) + (y + 3)³ = 0 ∴ -(x - 2)² - (x - 2)(y + 3) + (y + 3)² = 0[By (2)] ∴ (x - 2)² + (x - 2)(y + 3) - (y + 3)² = 0 ∴ (x² - 4x + 4) + (xy + 3x - 2y - 6) - (y² + 6y + 9) = 0 ∴ x² - 4x + 4 + xy + 3x - 2y - 6 - y² - 6y - 9 = 0

$$\therefore x^{2} + xy - y^{2} - x - 8y - 11 = 0$$

Exercise 4.3 | Q 2 | Page 127

Show that the equation $x^2 + 2xy + 2y^2 + 2x + 2y + 1 = 0$ does not represent a pair of lines.

Solution: Comparing the equation

 $x^{2} + 2xy + 2y^{2} + 2x + 2y + 1 = 0$ with $ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$, we get, a = 1, h = 1, b = 2, g = 1, f = 1, c = 1.

The given equation represents a pair of lines, if



$$D = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0 \text{ and } h^2 - ab \ge 0$$

Now, $D = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 1 \end{vmatrix}$

$$= 1(2 - 1) - 1(1 - 1) + 1(1 - 2)$$

$$= 1 - 0 - 1 = 0$$

and $h^2 - ab = (1)^2 - 1(2) = -1 < 0$

 \therefore given equation does not represent a pair of lines.

Exercise 4.3 | Q 3 | Page 127

Show that the equation $2x^2 - xy - 3y^2 - 6x + 19y - 20 = 0$ represents a pair of lines.

Solution: Comparing the equation

 $2x^2 - xy - 3y^2 - 6x + 19y - 20 = 0$

with $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, we get,

a = 2, h =
$$-\frac{1}{2}$$
, b = -3, f = $\frac{19}{2}$ and c = -20
∴ D = $\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}$

$$= \begin{vmatrix} 2 & -\frac{1}{2} & -3 \\ -\frac{1}{2} & -3 & \frac{19}{2} \\ -3 & \frac{19}{2} & -20 \end{vmatrix}$$

Taking $\frac{1}{2}$ common from each row, we get,



$$D = \frac{1}{8} \begin{vmatrix} 4 & -1 & -6 \\ -1 & -6 & 19 \\ -6 & 19 & -40 \end{vmatrix}$$

= $\frac{1}{8} [4(240 - 361) + 1(40 + 114) - 6(-19 - 36)]$
= $\frac{1}{8} [4(-121) + 154 - 6(-55)]$
= $\frac{11}{8} [4(-11) + 14 - 6(-5)]$
= $\frac{11}{8} (-44 + 14 + 30) = 0$
Also, h² - ab = $\left(-\frac{1}{2}\right)^2 - 2(-3) = \frac{1}{4} + 6 = \frac{25}{4} > 0$

 \therefore the given equation represents a pair of lines.

Exercise 4.3 | Q 4 | Page 127

Show that the equation $2x^2 + xy - y^2 + x + 4y - 3 = 0$ represents a pair of lines. Also, find the acute angle between them.

Solution: Comparing the equation

 $2x^{2} + xy - y^{2} + x + 4y - 3 = 0$ with $ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$ we get,

a = 2, h =
$$\frac{1}{2}$$
, b = -1, g = $\frac{1}{2}$, f = 2, c = -3.
∴ D = $\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = \begin{vmatrix} 2 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -1 & 2 \\ \frac{1}{2} & 2 & -3 \end{vmatrix}$
Taking $\frac{1}{2}$ common from each row, we get,



$$D = \frac{1}{8} \begin{vmatrix} 4 & 1 & 1 \\ 1 & -2 & 4 \\ 1 & 4 & -6 \end{vmatrix}$$
$$= \frac{1}{8} [4(12 - 16) - 1(-6 - 4) + 1(4 + 2)]$$
$$= \frac{1}{8} [4(-4) - 1(-10) + 1(6)]$$
$$= \frac{1}{8} (-16 + 10 + 6) = 0$$
Also, h² - ab = $\left(\frac{1}{2}\right)^2 - 2(-1) = \frac{1}{4} + 2 = \frac{9}{4} > 0$

 \therefore the given equation represents a pair of lines.

Let θ be the acute angle between the lines

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$= \left| \frac{2\sqrt{\left(\frac{1}{2}\right)^2 - 2(-1)}}{2 - 1} \right|$$
$$= \left| \frac{2\sqrt{\frac{1}{4} + 2}}{1} \right|$$
$$= 2 \times \frac{3}{2} = 3$$
$$\therefore \theta = \tan^{-1}(3)$$

Exercise 4.3 | Q 5.1 | Page 127



Find the separate equation of the line represented by the following equation:

$$(x - 2)^{2} - 3(x - 2)(y + 1) + 2(y + 1)^{2} = 0$$
Solution: $(x - 2)^{2} - 3(x - 2)(y + 1) + 2(y + 1)^{2} = 0$

$$\therefore (x - 2)^{2} - 2(x - 2)(y + 1) - (x - 2)(y + 1) + 2(y + 1)^{2} = 0$$

$$\therefore (x - 2)[(x - 2) - 2(y + 1)] - (y + 1)[(x - 2) - 2(y + 1)] = 0$$

$$\therefore (x - 2)(x - 2 - 2y - 2) - (y + 1)(x - 2 - 2y - 2) = 0$$

$$\therefore (x - 2)(x - 2y - 4) - (y + 1)(x - 2y - 4) = 0$$

$$\therefore (x - 2y - 4)(x - 2 - y - 1) = 0$$

$$\therefore (x - 2y - 4)(x - y - 3) = 0$$

$$\therefore \text{ the separate equations of the lines are}$$

$$x - 2y - 4 = 0 \text{ and } x - y - 3 = 0$$
Alternative Method:

 $(x - 2)^{2} - 3(x - 2)(y + 1) + 2(y + 1)^{2} = 0 \quad ...(1)$ Put x - 2 = X and y + 2 = Y \therefore (1) becomes, X² - 3XY + 2Y² = 0 \therefore X² - 2XY - XY + 2Y² = 0 \therefore X(X - 2Y) - Y(X - 2Y) = 0 \therefore (X - 2Y)(X - Y) = 0 \therefore the separate equations of the lines are X - 2Y = 0 and X - Y = 0 \therefore (x - 2) - 2(y + 1) = 0 and (x - 2) - (y + 1) = 0 \therefore x - 2y - 4 = 0 and x - y - 3 = 0

Exercise 4.3 | Q 5.2 | Page 127

Find the separate equation of the line represented by the following equation:

 $10(x + 1)^{2} + (x + 1)(y - 2) - 3(y - 2)^{2} = 0$ Solution: $10(x + 1)^{2} + (x + 1)(y - 2) - 3(y - 2)^{2} = 0$ (1) Put x + 1 = X and y - 2 = Y \therefore (1) becomes $10X^{2} + XY - 3Y^{2} = 0$



 $10X^{2} + 6XY - 5XY - 3Y^{2} = 0$ 2X(5X + 3Y) - Y(5X + 3Y) = 0 (2X - Y)(5X + 3Y) = 0 5X + 3Y = 0 and 2X - Y = 0 5X + 3Y = 0 5(x + 1) + 3(y - 2) = 0 5x + 5 + 3y - 6 = 0 $\therefore 5x + 3y - 1 = 0$ 2X - Y = 0 2(x + 1) - (y - 2) = 0 2x + 2 - y + 2 = 0 $\therefore 2x - y + 4 = 0$

Exercise 4.3 | Q 6.1 | Page 127

Find the value of k, if the following equations represent a pair of line:

 $3x^{2} + 10xy + 3y^{2} + 16y + k = 0$ **Solution:** Comparing the given equation with $ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$ we get, a = 3, h = 5, b = 3, g = 0, f = 8, c = k. Now, given equation represents a pair of lines. $\therefore abc + 2fgh - af^{2} - bg^{2} - ch^{2} = 0$ $\therefore (3)(3)(k) + 2(8)(0)(5) - 3(8)^{2} - 3(0)^{2} - k(5)^{2} = 0$ $\therefore 9k + 0 - 192 - 0 - 25k = 0$ $\therefore - 16k - 192 = 0$ $\therefore - 16k = 192$ $\therefore k = -12$

Exercise 4.3 | Q 6.2 | Page 127

Find the value of k, if the following equations represent a pair of line:

kxy + 10x + 6y + 4 = 0

Solution: Comparing the given equation with



$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$$

Now, given equation represents a pair of lines.

 $\therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0$

$$\therefore (0)(0)(4) + 2(3)(5)\left(\frac{k}{2}\right) - 0(3)^2 - 0(5)^2 - 4\left(\frac{k}{2}\right)^2 = 0$$

- $\therefore 0 + 15k 0 0 k^2 = 0$
- $\therefore 15k k^2 = 0$
- $\therefore k(k 15) = 0$
- \therefore k = 0 or k = 15
- If k = 0, then the given equation becomes
- 10x + 6y + 4 = 0 which does not represent a pair of lines.

∴ k ≠ 0

Hence, k = 15.

Exercise 4.3 | Q 6.3 | Page 127

Find the value of k, if the following equations represent a pair of line:

 $x^2 + 3xy + 2y^2 + x - y + k = 0$

Solution: Comparing the given equation with

 $ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$



we get, a = 1, h =
$$\frac{3}{2}$$
, b = 2, g = $\frac{1}{2}$, f = $-\frac{1}{2}$, c = k.

Now, given equation represents a pair of lines.

$$\therefore \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0$$
i.e.
$$\begin{vmatrix} 1 & \frac{3}{2} & \frac{1}{2} \\ \frac{3}{2} & 2 & -\frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & k \end{vmatrix} = 0$$

Taking out $\frac{1}{2}$ common from each row, we get,

 $\frac{1}{8}\begin{vmatrix} 2 & 3 & 1 \\ 3 & 4 & -1 \\ 1 & -1 & 2k \end{vmatrix} = 0$ $\therefore \begin{vmatrix} 2 & 3 & 1 \\ 3 & 4 & -1 \\ 1 & -1 & 2k \end{vmatrix} = 0$ $\therefore 2(8k - 1) - 3(6k + 1) + 1(-3 - 4) = 0$ $\therefore 16k - 2 - 18k - 3 - 7 = 0$ $\therefore -2k - 12 = 0$ $\therefore -2k = 12 \quad \therefore k = -6$

Exercise 4.3 | Q 7 | Page 128

Find p and q, if the equation $px^2 - 8xy + 3y^2 + 14x + 2y + q = 0$ represents a pair of perpendicular lines.

Solution: The given equation represents a pair of lines perpendicular to each other

: (coefficient of x^2) + (coefficient of y^2) = 0

$$\therefore p + 3 = 0 \quad \therefore p = -3$$

With this value of p, the given equation is



 $- 3x^{2} - 8xy + 3y^{2} + 14x + 2y + q = 0$ Comparing this equation with $ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$ a = -3, h = -4, b = 3, g = 7, f = 1 and c = q $\therefore D = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = \begin{vmatrix} -3 & -4 & 7 \\ -4 & 3 & 1 \\ 7 & 1 & g \end{vmatrix}$ = -3(3q - 1) + 4(-4q - 7) + 7(-4 - 21)= -9q + 3 - 16q - 28 - 175= -25q - 200= -25 (q + 8)

Since the given equation represents a pair of lines, D = 0

$$\therefore - 25(q+8) = 0$$

∴ q = - 8

Hence, p = -3 and q = -8.

Exercise 4.3 | Q 8 | Page 128

Find p and q, if the equation $2x^2 + 8xy + py^2 + qx + 2y - 15 = 0$ represents a pair of parallel lines.

Solution: The given equation is $2x^2 + 8xy + py^2 + qx + 2y - 15 = 0$ Comparing it with $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, we get, a = 2, h = 4, b = p, g = q/2, f = 1, c = -15Since the lines are parallel, $h^2 = ab$ $\therefore (4)^2 = 2p$ $\therefore p = 8$ Since the given equation represents a pair of lines



$$D = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0 \text{ where } b = p = 8$$

i.e.
$$\begin{vmatrix} 2 & 4 & \frac{q}{2} \\ 4 & 8 & 1 \\ \frac{q}{2} & 1 & -15 \end{vmatrix} = 0$$

i.e.
$$2(-120 - 1) - 4(-60 - \frac{q}{2}) + \frac{q}{2}(4 - 4q) = 0$$

i.e.
$$-242 + 240 + 2q + 2q - 2q^2 = 0$$

i.e.
$$-2q^2 + 4q - 2 = 0$$

i.e.
$$q^2 - 2q + 1 = 0$$

i.e.
$$(q - 1)^2 = 0$$

 $\therefore q = 1$
Hence, $p = 8$ and $q = 1$

Exercise 4.3 | Q 9 | Page 128

Equations of pairs of opposite sides of a parallelogram are $x^2 - 7x + 6 = 0$ and $y^2 - 14y + 40 = 0$. Find the joint equation of its diagonals.

Solution: Let ABCD be the parallelogram such that the combined equation of sides AB and CD is $x^2 - 7x + 6 = 0$ and the combined equation of sides BC and AD $y^2 - 14y + 40 = 0$

The separate equations of the lines represented by $x^2 - 7x + 6 = 0$, i.e. (x - 1)(x - 6) = 0are x - 1 = 0 and x - 6 = 0

Let equation of the side AB be x - 10 and equation of side CD be x - 6 = 0

The separate equations of the lines represented by $y^2 - 14y + 40 = 0$, i.e. (y - 4)(y - 10) = 0 are y - 4 = 0 and y - 10 = 0

Let equation of the side BC be y - 4 = 0 and equation of side AD be y - 10 = 0



Coordinates of the vertices of the parallelogram are A(1, 10), B(1, 4), C(6, 4) and D(6, 10)

 \therefore equation of the diagonal AC is

$$\frac{y-10}{x-1} = \frac{10-4}{1-6} = \frac{6}{-5}$$

$$\therefore 5y + 50 = 6x - 6$$

and equation of the diagonal BD is

$$\frac{y-4}{x-1} = \frac{4-10}{1-6} = \frac{-6}{-5} = \frac{6}{5}$$

$$\therefore 5y - 20 = 6x - 6$$

$$\therefore 6x - 5y + 14 = 0$$

Hence, the equations of the diagonals are 6x + 5y - 56 = 0 and 6x - 5y + 14 = 0

 \therefore the joint equation of the diagonals is

$$(6x + 5y - 56)(6x - 5y + 14) = 0$$

$$\therefore 36x^2 - 30xy + 84x + 30xy - 25y^2 + 70y - 336x + 280y - 784 = 0$$

$$\therefore 36x^2 - 25y^2 - 252x + 350y - 784 = 0$$

[Note: Answer in the textbook is incorrect]

Exercise 4.3 | Q 10 | Page 128


 $\triangle OAB$ is formed by the lines $x^2 - 4xy + y^2 = 0$ and the line AB. The equation of line AB is 2x + 3y - 1 = 0. Find the equation of the median of the triangle drawn from O. **Solution:**



Let D be the midpoint of seg AB where A is (x_1, y_1) and B is (x_2, y_2) .

Then D has coordinates $\left(rac{\mathrm{x}_1+\mathrm{x}_2}{2},rac{\mathrm{y}_1+\mathrm{y}_2}{2}
ight)$.

The joint (combined) equation of the lines OA and OB is $x^2 - 4xy + y^2 = 0$ and the equation of the line AB is 2x + 3y - 1 = 0

∴ points A and B satisfy the equations 2x + 3y - 1 = 0 and $x^2 - 4xy + y^2 = 0$ simultaneously.

We eliminate x from the above equations, i.e., put x = $\frac{1-3y}{2}$ in the equation x² - 4xy + y² = 0, we get,



$$\therefore \left(\frac{1-3y}{2}\right)^2 - 4\left(\frac{1-3y}{2}\right)y + y^2 = 0$$

$$\therefore (1-3y)^2 - 8(1-3y)y + 4y^2 = 0$$

$$\therefore 1 - 6y + 9y^2 - 8y + 24y^2 + 4y^2 = 0$$

$$\therefore 37y^2 - 14y + 1 = 0$$

The roots y_1 and y_2 of the above quadratic equation are the ycoordinates of the points A and B.

$$\therefore y_1 + y_2 = -\frac{b}{a} = \frac{14}{37}$$

$$\therefore y \text{-coordinate of } D = \frac{y_1 + y_2}{2} = \frac{7}{37}$$

Since D lies on the line AB, we can find the x-coordinate of D as

$$2x + 3\left(\frac{7}{37}\right) - 1 = 0$$

$$\therefore 2x = 1 - \frac{21}{37} = \frac{16}{37}$$

$$\therefore x = \frac{8}{37}$$

$$\therefore D \text{ is } \left(\frac{8}{37}, \frac{7}{37}\right)$$

$$\therefore \text{ equation of the median OD is } \frac{x}{8/37} = \frac{y}{7/37}$$

i.e. 7x - 8y = 0

Exercise 4.3 | Q 11 | Page 128



Find the coordinates of the points of intersection of the lines represented by $x^2 - y^2 - 2x$

+ 1 = 0
Solution: Consider,
$$x^2 - y^2 - 2x + 1 = 0$$

 $\therefore (x^2 - 2x + 1) - y^2 = 0$
 $\therefore (x - 1)^2 - y^2 = 0$
 $\therefore (x - 1 + y)(x - 1 - y) = 0$
 $\therefore (x + y - 1)(x - y - 1) = 0$
 \therefore separate equations of the lines are
 $x + y - 1 = 0$ and $x - y + 1 = 0$
To find the point of intersection of the lines, we have to solve
 $x + y - 1 = 0$...(1)
and $x - y + 1 = 0$...(2)
Adding (1) and (2), we get,
 $2x = 0$
 $\therefore x = 0$
Substituting $x = 0$ in (1), we get,
 $0 + y - 1 = 0$
 $\therefore y = 1$

 \therefore coordinates of the point of intersection of the lines are (0, 1).

[Note: Answer in the textbook is incorrect.]

MISCELLANEOUS EXERCISE 4 [PAGES 129 - 130]

Miscellaneous Exercise 4 | Q 1.01 | Page 129

Choose correct alternatives:

If the equation $4x^2 + hxy + y^2 = 0$ represents two coincident lines, then h = _____

- 1. ±2
- 2. ±3
- 3. ±4
- 4. ±5

Solution: If the equation $4x^2 + hxy + y^2 = 0$ represents two coincident lines, then $h = \pm 4$.



Miscellaneous Exercise 4 | Q 1.02 | Page 129

Choose correct alternatives:

If the lines represented by $kx^2 - 3xy + 6y^2 = 0$ are perpendicular to each other, then

1. k = 6

- 3. k = 3
- 4. k = 3

Solution: k = - 6

Miscellaneous Exercise 4 | Q 1.03 | Page 129

Choose correct alternatives:

Auxiliary equation of $2x^2 + 3xy - 9y^2 = 0$ is

- 1. $2m^2 + 3m 9 = 0$
- 2. $9m^2 3m 2 = 0$
- 3. $2m^2 3m + 9 = 0$
- 4. $-9m^2 3m + 2 = 0$

Solution: Auxiliary equation of $2x^2 + 3xy - 9y^2 = 0$ is $9m^2 - 3m - 2 = 0$.

Miscellaneous Exercise 4 | Q 1.04 | Page 129

Choose correct alternatives:

The difference between the slopes of the lines represented by $3x^2 - 4xy + y^2 = 0$ is 2

- 1. 2
- 2. 1
- 3. 3
- 4. 4

Solution: The difference between the slopes of the lines represented by $3x^2 - 4xy + y^2 =$

0 is <u>2.</u>

Miscellaneous Exercise 4 | Q 1.05 | Page 129

Choose correct alternatives:

If two lines $ax^2 + 2hxy + by^2 = 0$ make angles α and β with X-axis, then tan $(\alpha + \beta) =$

- 1. h/a+b
- 2. h/a-b
- 3. 2h/a+b



4. 2h/a-b

Solution:

If two lines $ax^2 + 2hxy + by^2 = 0$ make angles α and β with X-axis, then tan $(\alpha + \beta) = \frac{2h}{a - b}$

Explanation:

$$\begin{split} m_1 &= \tan \alpha, \, m_2 = \tan \beta \\ \therefore \, \tan \left(\alpha + \beta \right) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \cdot \tan \beta} \\ &= \frac{m_1 + m_2}{1 - m_1 m_2} = \frac{-2h/b}{1 - (a/b)} = \frac{2h}{a - b} \end{split}$$

Miscellaneous Exercise 4 | Q 1.06 | Page 129

Choose correct alternatives:

If the slope of one of the two lines given by $\frac{x^2}{a} + \frac{2xy}{h} + \frac{y^2}{b} = 0$ is twice that of the other, then ab : $h^2 =$ ____. 1. 1:2 2. 2:1 3. 8:9

4. 9:8

Solution:

If the slope of one of the two lines given by $\frac{x^2}{a} + \frac{2xy}{h} + \frac{y^2}{b} = 0$ is twice that of the other, then ab : $h^2 = \underline{9:8}$.

Explanation:

 m_1 + m_2 = $\frac{-2 \mathbf{b}}{\mathbf{h}}$ and $m_1 m_2$ = $\frac{\mathbf{b}}{\mathbf{a}}$



where $m_1 = 2m_2$

$$\therefore 2m_2 + m_2 = -\frac{2b}{h} \text{ and } 2m_2 \times m_2 = \frac{b}{a}$$
$$\therefore m_2 = \frac{-2b}{3h} \text{ and } m_2^2 = \frac{b}{2a}$$
$$\therefore \left(\frac{-2b}{3h}\right)^2 = \frac{b}{2a}$$
$$\therefore \frac{4b^2}{9h^2} = \frac{b}{2a}$$
$$\therefore \frac{4b^2}{9h^2} = \frac{b}{2a}$$
$$\therefore \frac{ab}{h^2} = \frac{9}{8}$$

Miscellaneous Exercise 4 | Q 1.07 | Page 130

Choose correct alternatives:

The joint equation of the lines through the origin and perpendicular to the pair of lines $3x^2 + 4xy - 5y^2 = 0$ is _____.

- 1. $5x^2 + 4xy 3y^2 = 0$
- 2. $3x^2 + 4xy 5y^2 = 0$
- 3. $3x^2 4xy + 5y^2 = 0$
- 4. $5x^2 + 4xy + 3y^2 = 0$

Solution: The joint equation of the lines through the origin and perpendicular to the pair of lines $3x^2 + 4xy - 5y^2 = 0$ is $5x^2 + 4xy - 3y^2 = 0$.

Miscellaneous Exercise 4 | Q 1.08 | Page 130

Choose correct alternatives:

If acute angle between lines $ax^2 + 2hxy + by^2 = 0$ is, $\pi/4$, then $4h^2 =$ _____.

- 1. $a^2 + 4ab + b^2$
- 2. $a^2 + 6ab + b^2$
- 3. (a + 2b)(a + 3b)
- 4. (a 2b)(2a + b)

Solution:



If acute angle between lines $ax^2 + 2hxy + by^2 = 0$ is, $\frac{\pi}{4}$, then $4h^2$

Miscellaneous Exercise 4 | Q 1.09 | Page 130

Choose correct alternatives:

If the equation $3x^2 - 8xy + qy^2 + 2x + 14y + p = 1$ represents a pair of perpendicular lines, then the values of p and q are respectively.

- 1. 3 and 7
- 2. 7 and 3
- 3. 3 and 7
- 4. 7 and 3

Solution: - 7 and - 3

Miscellaneous Exercise 4 | Q 1.1 | Page 130

Choose correct alternatives:

The area of triangle formed by the lines $x^2 + 4xy + y^2 = 0$ and x - y - 4 = 0 is

- 1. $4/\sqrt{3}$ sq units
- 2. $8/\sqrt{3}$ sq units
- 3. 16/ $\sqrt{3}$ sq units
- 4. $15/\sqrt{3}$ sq units

Solution: The area of triangle formed by the lines $x^2 + 4xy + y^2 = 0$ and x - y - 4 = 0

is 8/√3 sq units

Miscellaneous Exercise 4 | Q 1.11 | Page 130

Choose correct alternatives:

The combined equation of the coordinate axes is

- 1. x + y = 0
- 2. xy = k
- 3. xy = 0
- 4. x y = k

Solution: The combined equation of the coordinate axes is xy = 0.

Miscellaneous Exercise 4 | Q 1.12 | Page 130

Choose correct alternatives:

If $h^2 = ab$, then slopes of lines $ax^2 + 2hxy + by^2 = 0$ are in the ratio



- 1. 1:2
- 2. 2:1
- 3. 2:3
- 4. 1:1

Solution: If $h^2 = ab$, then slopes of lines $ax^2 + 2hxy + by^2 = 0$ are in the ratio <u>1:1.</u> **Hint:** If $h^2 = ab$, then lines are coincident. Therefore slopes of the lines are equal.

Miscellaneous Exercise 4 | Q 1.13 | Page 130

Choose correct alternatives:

If slope of one of the lines $ax^2 + 2hxy + by^2 = 0$ is 5 times the slope of the other, then $5h^2 =$ _____

- 1. ab
- 2. 2ab
- 3. 7ab
- 4. 9ab

Solution: If slope of one of the lines $ax^2 + 2hxy + by^2 = 0$ is 5 times the slope of the other, then $5h^2 = \underline{9ab}$.

Miscellaneous Exercise 4 | Q 1.14 | Page 130

Choose correct alternatives:

If distance between lines $(x - 2y)^2 + k(x - 2y) = 0$ is 3 units, then k =_____.

- 1. ±3
- 2. ± 5√5
- 3. 0
- 4. ±3√5

Solution: If distance between lines $(x - 2y)^2 + k(x - 2y) = 0$ is 3 units, then $k = \pm 3\sqrt{5}$

Explanation:

$$(x - 2y)^2 + k(x - 2y) = 0$$

$$\therefore (x - 2y)(x - 2y + k) = 0$$

 \therefore equations of the lines are x - 2y = 0 and x - 2y + k = 0 which are parallel to each other.



$$\therefore \left| \frac{\mathbf{k} - \mathbf{0}}{\sqrt{1 + 4}} \right| = 3$$

 \therefore k = $\pm 3\sqrt{5}$

MISCELLANEOUS EXERCISE 4 [PAGES 130 - 132]

Miscellaneous Exercise 4 | Q 1.01 | Page 130

Find the joint equation of the line:

x - y = 0 and x + y = 0

Solution: Find the joint equation of the line x - y = 0 and x + y = 0 is

(x - y)(x + y) = 0 $\therefore x^2 - y^2 = 0$

Miscellaneous Exercise 4 | Q 1.02 | Page 130

Find the joint equation of the line:

x + y - 3 = 0 and 2x + y - 1 = 0

Solution: Find the joint equation of the line x + y - 3 = 0 and 2x + y - 1 = 0

(x + y - 3)(2x + y - 1) = 0 $\therefore 2x^{2} + xy - x + 2xy + y^{2} - y - 6x - 3y + 3 = 0$ $\therefore 2x^{2} + 3xy + y^{2} - 7x - 4y + 3 = 0$

Miscellaneous Exercise 4 | Q 1.03 | Page 130

Find the joint equation of the line passing through the origin having slopes 2 and 3.

Solution: We know that the equation of the line passing through the origin and having slope m is y = mx. Equations of the lines passing through the origin and having slopes 2 and 3 are y = 2x and y = 3x respectively. i.e. their equations are

2x - y = 0 and 3x - y = 0 respectively.

 \therefore their joint equation is

(2x - y)(3x - y) = 0

 $\therefore 6x^2 - 2xy - 3xy + y^2 = 0$



$\therefore 6x^2 - 5xy + y^2 = 0$

Miscellaneous Exercise 4 | Q 1.04 | Page 130

Find the joint equation of the line passing through the origin and having inclinations 60° and 120°.

Solution: Slope of the line having inclination θ is tan θ .

Inclinations of the given lines are 60° and 120°

: their slopes are
$$m_1 = \tan 60^\circ = \sqrt{3}$$
 and

$$m_2 = \tan 120^\circ = \tan (180^\circ - 60^\circ)$$

$$= - \tan 60^\circ = -\sqrt{3}.$$

Since the lines pass through the origin, their equations are

y =
$$\sqrt{3}x$$
 and y = $-\sqrt{3}x$
i.e. $\sqrt{3}x$ - y = 0 and $\sqrt{3}x$ + y = 0

: the joint equation of these lines is

$$\left(\sqrt{3}x - y\right)\left(\sqrt{3}x + y\right) = 0$$

$$\therefore 3x^2 - y^2 = 0$$

Miscellaneous Exercise 4 | Q 1.05 | Page 130

Find the joint equation of the line passing through (1, 2) and parallel to the coordinate axes

Solution: Equations of the coordinate axes are x = 0 and y = 0

: the equations of the lines passing through (1, 2) and parallel to the coordinate axes are x = 1 and y = 2.

i.e. x - 1 = 0 and y - 2 = 0

 \therefore their combined equation is

$$(x - 1)(y - 2) = 0$$

$$\therefore x(y - 2) - 1(y - 2) = 0$$

$$\therefore xy - 2x - y + 2 = 0$$

Miscellaneous Exercise 4 | Q 1.06 | Page 130



Find the joint equation of the line passing through (3, 2) and parallel to the lines x = 2

and y = 3.

Solution: Equations of the lines passing through (3, 2) and parallel to the lines x = 2 and y = 3 are x = 3 and y = 2.

i.e. x - 3 = 0 and y - 2 = 0

 \therefore their joint equation is

(x - 3)(y - 2) = 0

 $\therefore xy - 2x - 3y + 6 = 0$

Miscellaneous Exercise 4 | Q 1.07 | Page 131

Find the joint equation of the line passing through (-1, 2) and perpendicular to x + 2y + 3 = 0 and 3x - 4y - 5 = 0

Solution: Let L_1 and L_2 be the lines passing through the origin and perpendicular to the lines x + 2y + 3 = 0 and 3x - 4y - 5 = 0 respectively.

Slopes of the lines x + 2y + 3 = 0 and 3x - 4y - 5 = 0 are $-\frac{1}{2}$ and $-\frac{3}{-4} = \frac{3}{4}$ respectively.

 \therefore slopes of the lines L_1 and L_2 are 2 and $\frac{-4}{3}$ respectively.

Since the lines L_1 and L_2 pass through the point (-1, 2), their equations are

 $\therefore (y - y_1) = m(x - x_1)$

$$(y - 2) = 2(x + 1)$$



⇒ 2x - y + 4 = 0 and

$$\therefore (y-2) = \left(\frac{-4}{3}\right)(x+1)$$

 \Rightarrow 3y - 6 = - 4x - 4

 $\Rightarrow 4x + 3y - 6 + 4 = 0$

 \Rightarrow 4x + 3y - 2 = 0

their combined equation is

- $\therefore (2x y + 4)(4x + 3y 2) = 0$
- $\therefore 8x^2 + 6xy 4x 4xy 3y^2 + 2y + 16x + 12y 8 = 0$
- $\therefore 8x^2 + 2xy + 12x 3y^2 + 14y 8 = 0$

Miscellaneous Exercise 4 | Q 1.08 | Page 131

Find the joint equation of the line passing through the origin and having slopes 1 + $\sqrt{3}$ and 1 - $\sqrt{3}$

Solution:

Let I₁ and I₂ be the two lines. Slopes of I₁ is 1 + $\sqrt{3}$ and that of I₂ is 1 - $\sqrt{3}$

Therefore the equation of a line (I_1) passing through the origin and having slope is

$$y = (1 + \sqrt{3})x$$
$$\therefore (1 + \sqrt{3})x - y = 0 \quad \dots (1)$$

Similarly, the equation of the line (I2) passing through the origin and having slope is

$$y = (1 - \sqrt{3})x$$



$$\therefore \left(1 - \sqrt{3}\right) \mathbf{x} - \mathbf{y} = 0 \quad \dots (2)$$

From (1) and (2) the required combined equation is

$$\begin{split} &\left[\left(1 + \sqrt{3} \right) x - y \right] \left[\left(1 - \sqrt{3} \right) x - y \right] = 0 \\ &\therefore \left(1 + \sqrt{3} \right) x \left[\left(1 - \sqrt{3} \right) x - y \right] - y \left[\left(1 - \sqrt{3} \right) x - y \right] = 0 \\ &\therefore \left(1 - \sqrt{3} \right) \left(1 + \sqrt{3} \right) x^2 - \left(1 + \sqrt{3} \right) xy - \left(1 - \sqrt{3} \right) xy + y^2 = 0 \\ &\therefore \left((1)^2 - \left(\sqrt{3} \right)^2 \right) x^2 - \left[\left(1 + \sqrt{3} \right) + \left(1 - \sqrt{3} \right) \right] xy + y^2 = 0 \\ &\therefore (1 - 3) x^2 - 2xy + y^2 = 0 \\ &\therefore -2x^2 - 2xy + y^2 = 0 \\ &\therefore 2x^2 + 2xy - y^2 = 0 \end{split}$$

This is the required combined equation.

Miscellaneous Exercise 4 | Q 1.09 | Page 131

Find the joint equation of the line which are at a distance of 9 units from the Y-axis.

Solution: Equations of the lines, which are parallel to the Y-axis and at a distance of 9 units from it, are x = 9 and x = -9

i.e. x - 9 = 0 and x + 9 = 0



 \therefore their combined equation is



(x - 9)(x + 9) = 0 $\therefore x^2 - 81 = 0$

Miscellaneous Exercise 4 | Q 1.1 | Page 131

Find the joint equation of the line passing through the point (3, 2), one of which is parallel to the line x - 2y = 2, and other is perpendicular to the line y = 3.

Solution:

Let L_1 be the line passes through (3, 2) and parallel to the line x -

2y = 2 whose slope is
$$\frac{-1}{-2} = \frac{1}{2}$$

∴ slope of the line L₁ is $\frac{1}{2}$

 \therefore equation of the line L₂ is

$$y - 2 = \frac{1}{2}(x - 3)$$

$$\therefore x - 2y + 1 = 0$$

Let L_2 be the line passes through (3, 2) and perpendicular to the line y = 3.

 \therefore equation of the line L₂ is of the form x = a. Since L₂ passes through (3, 2), 3 = a.

 \therefore equation of the line L₂ is x = 3, i.e. x - 3 = 0

Hence, the equations of the required lines are

$$x - 2y + 1 = 0$$
 and $x - 3 = 0$

 \therefore their joint equation is

$$(x - 2y + 1)(x - 3) = 0$$

$$\therefore x^2 - 2xy + x - 3x + 6y - 3 = 0$$

$$\therefore x^2 - 2xy - 2x + 6y - 3 = 0$$

Miscellaneous Exercise 4 | Q 1.11 | Page 131

Find the joint equation of the line passing through the origin and perpendicular to the lines x + 2y = 19 and 3x + y = 18



Solution: Let L_1 and L_2 be the lines passing through the origin and perpendicular to the lines x + 2y = 19 and 3x + y = 18 respectively.

Slopes of the lines x + 2y = 19 and 3x + y = 18 are -1/2 and -3/1 = -3 respectively.

 \therefore slopes of the lines L₁ and L₂ are 2 and 1/3 respectively.

Since the lines L1 and L2 pass through the origin, their equations are

y = 2x and y = 1/3 x

i.e. 2x - y = 0 and x - 3y = 0

 \therefore their combined equation is

(2x - y)(x - 3y) = 0

 $\therefore 2x^2 - 6xy - xy + 3y^2 = 0$

 $\therefore 2x^2 - 7xy + 3y^2 = 0$

Miscellaneous Exercise 4 | Q 2.1 | Page 131

Show that the following equations represents a pair of line:

$$x^2 + 2xy - y^2 = 0$$

Solution: Comparing the equation $x^2 + 2xy - y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,

a = 1, 2h = 2 i,e, h = 1, and b = - 1 ∴ h² - ab = (1)² - 1(- 1) = 1 + 2 = 2 > 0

Since the equation $x^2 + 2xy - y^2 = 0$ is a homogeneous equation of second degree and $h^2 - ab > 0$, the given equation represents a pair of lines which are real and distinct.

Miscellaneous Exercise 4 | Q 2.2 | Page 131

Show that the following equations represents a pair of line:

$$4x^2 + 4xy + y^2 = 0$$

Solution: Comparing the equation $4x^2 + 4xy + y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,

$$a = 4$$
, $2h = 4$ i,e, $h = 2$, and $b = 1$

$$h^2 - ab = (2)^2 - 4(1) = 4 - 4 = 0$$

Since the equation $4x^2 + 4xy + y^2 = 0$ is a homogeneous equation of second degree and h^2 - ab = 0, the given equation represents a pair of lines which are real and coincident.



Miscellaneous Exercise 4 | Q 2.3 | Page 131

Show that the following equations represent a pair of line:

$$x^2 - y^2 = 0$$

Solution: Comparing the equation $x^2 - y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,

a = 1, 2h = 0 i,e, h = 0, and b = -1

$$\therefore$$
 h² - ab = (0)² - 1 (- 1) = 0 + 1 = 1 > 0

Since the equation $x^2 - y^2 = 0$ is a homogeneous equation of second degree and $h^2 - ab > 0$, the given equation represents a pair of lines which are real and distinct.

Miscellaneous Exercise 4 | Q 2.4 | Page 131

Show that the following equations represent a pair of line:

 $x^2 + 7xy - 2y^2 = 0$

Solution: Comparing the equation
$$x^2 + 7xy - 2y^2 = 0$$
 with $ax^2 + 2hxy + by^2 = 0$, we get,
 $a = 1, 2h = 7$ i.e., $h = \frac{7}{2}$, and $b = -2$
 $\therefore h^2 - ab = \left(\frac{7}{2}\right)^2 - 1$ (-2)
 $= \frac{49}{4} + 2$
 $= \frac{57}{4}$ i.e. 14.25 = 14 > 0

Since the equation $x^2 + 7xy - 2y^2 = 0$ is a homogeneous equation of second degree and $h^2 - ab > 0$, the given equation represents a pair of lines which are real and distinct. **Miscellaneous Exercise 4 | Q 2.5 | Page 131**

Show that the following equations represent a pair of line:

$$x^2 - 2\sqrt{3}xy - y^2 = 0$$

Solution:



Comparing the equation $x^2 - 2\sqrt{3}xy - y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, $a = 1, 2h = -2\sqrt{3}$ i,e, $h = \sqrt{3}$, and b = 1 $\therefore h^2 - ab = (\sqrt{3})^2 - 1$ (1) = 3 - 1 = 2 > 0

Since the equation $x^2 - 2\sqrt{3}xy - y^2 = 0$ is a homogeneous equation of second degree and h^2 - ab > 0, the given equation represents a pair of lines which are real and distinct.

Miscellaneous Exercise 4 | Q 3.1 | Page 131

Find the separate equation of the line represented by the following equation:

$$6x^{2} - 5xy - 6y^{2} = 0$$

Solution: $6x^{2} - 5xy - 6y^{2} = 0$
 $\therefore 6x^{2} - 9xy + 4xy - 6y^{2} = 0$
 $\therefore 3x (2x - 3y) + 2y(2x - 3y) = 0$
 $\therefore (2x - 3y)(3x + 2y) = 0$
the separate equations of the lines are

2x - 3y = 0 and 3x + 2y = 0.

Miscellaneous Exercise 4 | Q 3.2 | Page 131

Find the separate equation of the line represented by the following equation:

$$x^{2} - 4y^{2} = 0$$

Solution: $x^{2} - 4y^{2} = 0$
 $\therefore x^{2} - (2y)^{2} = 0$
 $\therefore (x - 2y)(x + 2y) = 0$
the separate equations of the lines are
 $x - 2y = 0$ and $x + 2y = 0$

Miscellaneous Exercise 4 | Q 3.3 | Page 131



Find the separate equation of the line represented by the following equation:

 $3x^2 - y^2 = 0$

Solution:

$$3x^{2} - y^{2} = 0$$

$$\therefore \left(\sqrt{3}x\right)^{2} - y^{2} = 0$$

$$\therefore \left(\sqrt{3}x - y\right)\left(\sqrt{3}x + y\right) = 0$$

the separate equations of the lines are

$$\sqrt{3}x-y=0$$
 and $\sqrt{3}x+y=0$

Miscellaneous Exercise 4 | Q 3.4 | Page 131

Find the separate equation of the line represented by the following equation:

$$2x^{2} + 2xy - y^{2} = 0$$
Solution: $2x^{2} + 2xy - y^{2} = 0$
The auxiliary equation is $-m^{2} + 2m + 2 = 0$

$$\therefore m^{2} - 2m - 2 = 0$$

$$\therefore m = \frac{2 \pm \sqrt{(-2)^{2} - 4(1)(-2)}}{2 \times 1}$$

$$= \frac{2 \pm \sqrt{4 + 8}}{2}$$

$$= \frac{2 \pm 2\sqrt{3}}{2}$$

$$= 1 \pm \sqrt{3}$$

$$\therefore m_{1} = 1 + \sqrt{3} \text{ and } m_{2} = 1 - \sqrt{3} \text{ are the slopes of the lines.}$$

∴ their separate equations are



$$y = m_1 x$$
 and $y = m_2 x$

i.e.
$$y = (1 + \sqrt{3})x$$
 and $y = (1 - \sqrt{3})x$
i.e. $(\sqrt{3} + 1)x - y = 0$ and $(\sqrt{3} - 1)x + y = 0$

Miscellaneous Exercise 4 | Q 4.1 | Page 131

Find the joint equation of the pair of a line through the origin and perpendicular to the lines given by

$$x^2 + 4xy - 5y^2 = 0$$

Solution: Comparing the equation $x^2 + 4xy - 5y^2 = 0$ with $ax_2 + 2hxy + by^2 = 0$, we get,

a = 1, 2h = 4, b = -5

Let m_1 and m_2 be the slopes of the lines represented by $x^2 + 4xy - 5y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{4}{5}$$
 and $m_1m_2 = \frac{a}{b} = \frac{-1}{5}$...(1)

Now, required lines are perpendicular to these lines

$$\therefore$$
 their slopes are $\displaystyle\frac{-1}{m_1}$ and $\displaystyle\frac{1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$\mathsf{y} = \frac{-1}{m_1} \mathbf{x} \text{ and } \mathsf{y} = \frac{-1}{m_2} \mathbf{x}$$

i.e. $m_1y = -x$ and $m_2y = -x$ i.e. $x + m_1y = 0$ and $x + m_2y = 0$ \therefore their combined equation is $(x + m_1y)(x + m_2y) = 0$ $\therefore x^2 + (m_1 + m_2)xy + m_1m_2y^2 = 0$ $\therefore x^2 + \frac{4}{5}xy - \frac{1}{5}y^2 = 0$ [By(1)] $\therefore 5x^2 + 4xy - y^2 = 0$



Miscellaneous Exercise 4 | Q 4.2 | Page 131

Find the joint equation of the pair of a line through the origin and perpendicular to the lines given by

$$2x^2 - 3xy - 9y^2 = 0$$

Solution: Comparing the equation $2x^2 - 3xy - 9y^2 = 0$ with $ax_2 + 2hxy + by^2 = 0$, we get,

a = 2, 2h = - 3, b = - 9

Let m_1 and m_2 be the slopes of the lines represented by $2x^2 - 3xy - 9y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = -\frac{3}{9}$$
 and $m_1m_2 = \frac{a}{b} = -\frac{2}{9}$...(1)

Now, required lines are perpendicular to these lines

$$\therefore$$
 their slopes are $\displaystyle\frac{-1}{m_1}$ and $\displaystyle-\frac{1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$\mathsf{y} = rac{-1}{\mathrm{m}_1} \mathsf{x} ext{ and } \mathsf{y} = rac{-1}{\mathrm{m}_2} \mathsf{x}$$

i.e. $m_1y = -x$ and $m_2y = -x$

i.e.
$$x + m_1 y = 0$$
 and $x + m_2 y = 0$

 \div their combined equation is

$$(x + m_1 y)(x + m_2 y) = 0$$

$$\therefore x^2 + (m_1 + m_2)xy + m_1m_2y^2 = 0$$

$$\therefore \mathbf{x}^2 + \left(-\frac{3}{9}\right)\mathbf{x}\mathbf{y} + \left(-\frac{2}{9}\right)\mathbf{y}^2 = 0 \quad \dots [\mathsf{By}(1)]$$
$$\therefore 9\mathbf{x}^2 - 3\mathbf{x}\mathbf{y} - 2\mathbf{y}^2 = 0$$

Miscellaneous Exercise 4 | Q 4.3 | Page 131

Find the joint equation of the pair of a line through the origin and perpendicular to the lines given by

$$x^2 + xy - y^2 = 0$$

Solution: Comparing the equation $x^2 + xy - y^2 = 0$ with $ax_2 + 2hxy + by^2 = 0$, we get, a = 1, 2h = 1, b = -1



Let m_1 and m_2 be the slopes of the lines represented by $x^2 + xy - y^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{-1}{-1} = 1 \text{ and } m_1m_2 = \frac{a}{b} = \frac{1}{-1} = -1 \quad ...(1)$$

Now, required lines are perpendicular to these lines

$$\therefore$$
 their slopes are $\displaystyle\frac{-1}{m_1}$ and $\displaystyle-\frac{1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$\mathsf{y} = \frac{-1}{m_1} \mathbf{x} \text{ and } \mathsf{y} = \frac{-1}{m_2} \mathbf{x}$$

i.e. $m_1y = -x$ and $m_2y = -x$

i.e.
$$x + m_1 y = 0$$
 and $x + m_2 y = 0$

 \div their combined equation is

$$(x + m_1 y)(x + m_2 y) = 0$$

$$\therefore x^2 + (m_1 + m_2)xy + m_1m_2y^2 = 0$$

$$\therefore x^{2} + 1xy + (-1)y^{2} = 0$$
[By(1)]

$$\therefore x^2 + xy - y^2 = 0$$

Miscellaneous Exercise 4 | Q 5.1 | Page 131

Find k, if the sum of the slopes of the lines given by $3x^2 + kxy - y^2 = 0$ is zero. **Solution:** Comparing the equation $3x^2 + kxy - y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = 3, 2h = k, b = -1

Let m_1 and m_2 be the slopes of the lines represented by $3x^2 + kxy - y^2 = 0$.

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{-k}{-1} = k$$

Now, m₁ + m₂ = 0 ...(Given)

Miscellaneous Exercise 4 | Q 5.2 | Page 131



Find k, if the sum of the slopes of the lines given by $x^2 + kxy - 3y^2 = 0$ is equal to their product.

Solution: Comparing the equation $x^2 + kxy - 3y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = 1, 2h = k, b = - 3

Let m_1 and m_2 be the slopes of the lines represented by $x^2 + kxy - 3y^2 = 0$.

$$\therefore m_1 + m_2 = \frac{-2h}{b} = \frac{-k}{-3} = \frac{k}{3}$$
$$m_1 m_2 = \frac{a}{b} = \frac{1}{-3} = \frac{-1}{3}$$

Now, $m_1 + m_2 = m_1 m_2$...(Given)

$$\therefore \frac{\mathbf{k}}{3} = \frac{-1}{3}$$

∴ k = - 1.

Miscellaneous Exercise 4 | Q 5.3 | Page 131

Find k, if the slope of one of the lines given by $3x^2 - 4xy + ky^2 = 0$ is 1.

Solution: The auxiliary equation of the lines given by $3x^2 - 4xy + ky^2 = 0$ is $km^2 - 4m + 3 = 0$

Given, slope of one of the lines is 1.

 \therefore m = 1 is the root of the auxiliary equation km² - 4m + 3 = 0

$$\therefore k(1)^2 - 4(1) + 3 = 0$$

 $\therefore k - 4 + 3 = 0$

∴ k = 1

Miscellaneous Exercise 4 | Q 5.4 | Page 131

Find k, if one of the lines given by $3x^2 - kxy + 5y^2 = 0$ is perpendicular to the line 5x + 3y = 0.

Solution:



The auxiliary equation of the lines given by $3x^2 - kxy + 5y^2 = 0$ is $5m^2 - km + 3 = 0$ Now, one line is perpendicular to the line 5x + 3y = 0, whose slope is $-\frac{5}{3}$

$$\therefore \text{ slope of that line = m = } \frac{3}{5}$$

$$\therefore m = \frac{3}{5} \text{ is the root of the auxiliary equation } 5m^2 - km + 3 = 0$$

$$\therefore 5\left(\frac{3}{5}\right)^2 - k\left(\frac{3}{5}\right) + 3 = 0$$

$$\therefore \frac{9}{5} - \frac{3k}{5} + 3 = 0$$

$$\therefore 9 - 3k + 15 = 0$$

$$\therefore 3k = 24$$

$$\therefore k = 8$$

Miscellaneous Exercise 4 | Q 5.5 | Page 131

Find k if the slope of one of the lines given by $3x^2 + 4xy + ky^2 = 0$ is three times the other.

Solution: $3x^2 + 4xy + ky^2 = 0$ \therefore divide by x^2 $\frac{3x^2}{x^2} + \frac{4xy}{x^2} + \frac{ky^2}{x^2} = 0$ $3 + \frac{4y}{x} + \frac{ky^2}{x^2} = 0$ (1) \therefore y = mx \therefore $\frac{y}{x} = m$ Put $\frac{y}{x} = m$ in equation (1)

Comparing the equation $km^2 + 4m + 3 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,



a = k, 2h = 4, b = 3 $m_1 = 3m_2$...(given condition) $m_1 + m_2 = \frac{-2h}{r} = -\frac{4}{r}$ $m_1m_2 = \frac{a}{b} = \frac{3}{b}$ $m_1 + m_2 = -\frac{4}{r}$ $4m_2 = -\frac{4}{r}$ (m₁ = 3m₂) $m_2 = -\frac{1}{k}$ $m_1m_2 = \frac{3}{\nu}$ $3m_2^2 = \frac{3}{r}$ (m₁ = 3m₂) $3\left(-\frac{1}{k}\right)^2 = \frac{3}{k}$ (m₂ = $-\frac{1}{k}$) $\frac{1}{k^2} = \frac{1}{k}$ $k^2 = k$ k = 1 or k = 0

Miscellaneous Exercise 4 | Q 5.6 | Page 131

Find k, if the slopes of lines given by $kx^2 + 5xy + y^2 = 0$ differ by 1. Solution: Comparing the equation $kx^2 + 5xy + y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,



a = k, 2h = 5 i.e. h =
$$\frac{5}{2}$$

m₁ + m₂ = $\frac{-2h}{b} = -\frac{5}{1} = -5$
and m₁m₂ = $\frac{a}{b} = \frac{k}{1} = k$

the slope of the line differ by $(m_1 - m_2) = 1$ (1) $\therefore (m_1 - m_2)^2 - (m_1 + m_2)^2 - 4m_1m_2$

$$(m_1 - m_2)^2 = (-5)^2 - 4(k)$$

$$(m_1 - m_2)^2 = 25 - 4k$$

$$1 = 25 - 4k \qquad \dots [By(1)]$$

$$4k = 24$$

$$k = 6$$

Miscellaneous Exercise 4 | Q 5.7 | Page 131

Find k, if one of the lines given by $6x^2 + kxy + y^2 = 0$ is 2x + y = 0.

Solution: The auxiliary equation of the lines represented by $6x^2 + kxy + y^2 = 0$ is $m^2 + km + 6 = 0$

Since one of the line is 2x + y = 0 whose slope is m = -2.

 \therefore m = - 2 is the root of the auxiliary equation m² + km + 6 = 0.

$$\therefore (-2)^2 + k(-2) + 6 = 0$$

- :: 4 2k + 6 = 0
- ∴ 2k = 10
- ∴ k = 5

Miscellaneous Exercise 4 | Q 6 | Page 131

Find the joint equation of the pair of lines which bisect angles between the lines given by $x^2 + 3xy + 2y^2 = 0$ **Solution:** $x^2 + 3xy + 2y^2 = 0$ $\therefore x^2 + 2xy + xy + 2y^2 = 0$ $\therefore x(x + 2y) + y(x + 2y) = 0$



 $\therefore (x+2y)(x+y) = 0$

: separate equations of the lines represented by $x^2 + 3xy + 2y^2 = 0$ are x + 2y = 0 and x + y = 0

Let P (x, y) be any point on one of the angle bisector. Since the points on the angle bisectors are equidistant from both the lines,



the distance of P(x, y) from the line x + 2y = 0= the distance of P(x, y) from the line x + y = 0

$$\therefore \left| \frac{\mathbf{x} + 2\mathbf{y}}{\sqrt{1+4}} \right| = \left| \frac{\mathbf{x} + \mathbf{y}}{\sqrt{1+1}} \right|$$
$$\therefore \frac{(\mathbf{x} + 2\mathbf{y})^2}{5} = \frac{(\mathbf{x} + \mathbf{y})^2}{2}$$

 $\therefore 2(x + 2y)^{2} = 5(x + y)^{2}$ $\therefore 2(x^{2} + 4xy + 4y^{2}) = 5(x^{2} + 2xy + y^{2})$ $\therefore 2x^{2} + 8xy + 8y^{2} = 5x^{2} + 10xy + 5y^{2}$ $\therefore 3x^{2} + 2xy - 3y^{2} = 0$

This is the required joint equation of the lines which bisect the angles between the lines represented by $x^2 + 3xy + 2y^2 = 0$

Miscellaneous Exercise 4 | Q 7 | Page 131

Find the joint equation of the pair of lines through the origin and making an equilateral triangle with the line x = 3.

Solution:





Let OA and OB be the lines through the origin making an angle of 60° with the line x = 3.

 \therefore OA and OB make an angle of 30° and 150° with the positive direction of X-axis.

 \therefore slope of OA = tan 30° = $\frac{1}{\sqrt{3}}$ \therefore equation of the line OA is y = $\frac{1}{\sqrt{3}}$ x $\therefore \sqrt{3}y = x$ $\therefore x - \sqrt{y} = 0$ Slope of OB = tan 150° = tan (180° - 30°) $= - \tan 30^\circ = -\frac{1}{\sqrt{3}}$ \therefore equation of the line OB is y = $\frac{-1}{\sqrt{3}}$ x $\therefore \sqrt{3}y = -x$ $\therefore x + \sqrt{3}y = 0$... required combined equation of the lines is $\left(\mathbf{x}-\sqrt{3}\mathbf{y}\right)\left(\mathbf{x}+\sqrt{3}\mathbf{y}\right)=0$ i.e. $x^2 - 3y^2 = 0$



Miscellaneous Exercise 4 | Q 8 | Page 131

Show that the lines $x^2 - 4xy + y^2 = 0$ and x + y = 10 contain the sides of an equilateral triangle. Find the area of the triangle.

Solution: We find the joint equation of the pair of lines OA and OB through origin, each making an angle of 60° with x + y = 10 whose slope is - 1.

Let OA(or OB) has slope m.

 \therefore its equation is y - mx(1)

Also, tan 60° =
$$\left| \frac{m - (-1)}{1 + m(-1)} \right|$$

$$\therefore \sqrt{3} = \left| \frac{m + 1}{1 - m} \right|$$

Squaring both sides, we get,

$$3 = \frac{(m+1)^2}{(1-m)^2}$$

$$\therefore 3(1 - 2m + m^2) = m^2 + 2m + 1$$

$$\therefore 3 - 6m + 3m^2 = m^2 + 2m + 1$$

$$\therefore 2m^2 - 8m + 2 = 0$$

$$\therefore m^2 - 4m + 1 = 0$$

$$\therefore \left(\frac{y}{x}\right) - 4\left(\frac{y}{x}\right) + 1 = 0 \quad ...[By(1)]$$

$$\therefore y^2 - 4xy + x^2 = 0$$

 $\therefore x^2 - 4xy + y^2 = 0$ is the joint equation of the two lines through the origin each making an angle of 60° with x + y = 10

 $\therefore x^2 - 4xy + y^2 = 0$ and x + y = 10 form a triangle OAB which is equilateral.

Let seg OM perpendicular line AB whose question is x + y = 10



$$\therefore \text{ OM} = \left| \frac{-10}{\sqrt{1+1}} \right| = 5\sqrt{2}$$

$$\therefore \text{ area of equilateral } \Delta \text{ OAB} = \frac{(\text{OM})^2}{\sqrt{3}} = \frac{\left(5\sqrt{2}\right)^2}{\sqrt{3}}$$

$$= \frac{50}{\sqrt{3}} \text{ sq units.}$$

Miscellaneous Exercise 4 | Q 9 | Page 131

If the slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is three times the other, prove that $3h^2 = 4ab$.

Solution: Let m_1 and m_2 be the slopes of the lines represented by $ax^2 + 2hxy + by^2 = 0$ We are given that $m_2 = 3m_1$

 $\therefore m_{1} + 3m_{1} = -\frac{2h}{b}$ $\therefore 4m_{1} = -\frac{2h}{b}$ $\therefore m_{1} = -\frac{h}{2b}$ Also, $m_{1}(3m_{1}) = \frac{a}{b}$ $\therefore 3m_{1}^{2} = \frac{a}{b}$ $\therefore 3\left(-\frac{h}{2b}\right)^{2} = \frac{a}{b} \quad[By (1)]$ $\therefore \frac{3h^{2}}{4b^{2}} = \frac{a}{b}$ $\therefore 3h^{2} = 4ab, as b \neq 0$

Miscellaneous Exercise 4 | Q 10 | Page 132



Find the combined equation of bisectors of angles between the lines represented by $5x^2 + 6xy - y^2 = 0$.

Solution: Comparing the equation $5x^2 + 6xy - y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get,

a = 5, 2h = 6, b = -1

Let m_1 and m_2 be the slopes of the lines represented by $5x^2 + 6xy - y^2 = 0$.

$$\therefore m1 + m2 = \frac{-2h}{b} = \frac{-6}{-1} = 6 \text{ and } m_1m_2 = \frac{a}{b} = \frac{5}{-1} = -5$$
 ...(1)

The separate equations of the lines are

 $y = m_1 x$ and $y = m_2 x$, where $m_1 \neq m_2$

i.e. $m_1x - y = 0$ and $m_2x - y = 0$.

Let P(x, y) be any point on one of the bisector of the angles between the lines.

: the distance of P from the line $m_1x - y = 0$ is equal to the distance of P from the line $m_2x - y = 0$.

$$\therefore \left| rac{\mathrm{m_1x} - \mathrm{y}}{\sqrt{\mathrm{m_1^2} + 1}}
ight| = \left| rac{\mathrm{m_2x} - \mathrm{y}}{\sqrt{\mathrm{m_2^2} + 1}}
ight|$$

Squaring both sides, we get,

$$\begin{split} & \frac{(m_1x-y)^2}{m_1^2+1} = \frac{m_2x-y}{m_2^2+1} \\ & \therefore \ (m_2^2+1)(m_1x-y)^2 = (m_1^2+1)(m_2x-y) \\ & \therefore \ (m_2^2+1)(m_1^2x^2-2m_1xy+y^2) = (m_1^2+1)(m_2^2x^2-2m_2xy+y^2) \\ & \therefore \\ & m_1^2m_2^2x^2-2m_1m_2^2y^2xy+m_2^2y^2+m_1^2x^2-2m_1xy+y^2 = m_1^2m_2^2x^2-2m_1^2m_2xy+m_1^2y^2+m_2^2x^2-2m_2xy+y^2 \\ & \therefore \ (m_1^2-m_2^2)x^2+2m_1m_2(m_1-m_2)xy-2(m_1-m_2)xy-(m_1^2-m_2^2)y^2 = 0 \\ & \text{Dividing throughout by } m_1-m_2 \ (\neq 0), \text{ we get,} \\ & (m_1+m_2)x^2+2m_1m_2xy-2xy-(m_1+m_2)y^2 = 0 \end{split}$$



 $\therefore 6x^{2} - 10xy - 2xy - 6y^{2} = 0 \qquad ...[By (1)]$ $\therefore 6x^{2} - 12xy - 6y^{2} = 0$ $\therefore x^{2} - 2xy - y^{2} = 0$

This is the joint equation of the bisectors of the angles between the lines represented by $5x^2 + 6xy - y^2 = 0$.

Miscellaneous Exercise 4 | Q 11 | Page 132

Find an if the sum of the slope of lines represented by $ax^2 + 8xy + 5y^2 = 0$ is twice their product.

Solution: Comparing the equation $ax^2 + 8xy + 5y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get, a = a, 2h = 8, b = 5

Let m_1 and m_2 be the slopes of the lines represented by $ax^2 + 8xy + 5y^2 = 0$.

$$\therefore m_1 + m_2 = \frac{-2h}{b} = -\frac{8}{5}$$

and $m_1m_2 = \frac{a}{b} = \frac{a}{5}$

Now, $(m_1 + m_2) = 2(m_1m_2)$

$$-\frac{8}{5} = 2\left(\frac{a}{5}\right)$$

a = 4

Miscellaneous Exercise 4 | Q 12 | Page 132

If the line 4x - 5y = 0 coincides with one of the lines given by $ax^2 + 2hxy + by^2 = 0$, then show that 25a + 40h + 16b = 0

Solution: The auxiliary equation of the lines represented by $ax^2 + 2hxy + by^2 = 0$ is $bm^2 + 2hm + a = 0$.

Given that 4x - 5y = 0 is one of the lines represented by $ax^2 + 2hxy + by^2 = 0$.



The slope of the line 4x - 5y = 0 is $\frac{-4}{-5} = \frac{4}{5}$

 $\therefore m = \frac{4}{5} \text{ is a root of the auxiliary equation } bm^{2} + 2hm + a = 0.$ $\therefore b\left(\frac{4}{5}\right)^{2} + 2h\left(\frac{4}{5}\right) + a = 0$ $\therefore \frac{16b}{25} + \frac{8h}{5} + a = 0$ $\therefore 16b + 40h + 25a = 0 \text{ i.e.}$

∴ 25a + 40h + 16b = 0

Miscellaneous Exercise 4 | Q 13.1 | Page 132

Show that the following equation represents a pair of line. Find the acute angle between them:

 $9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0$

Solution: Comparing this equation with

 $ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$, we get,

a = 9, h = -3, b = 1, g = 9, f = -3 and c = 8

$$\therefore D = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}$$
$$= \begin{vmatrix} 9 & -3 & 9 \\ -3 & 1 & -3 \\ 9 & -3 & 8 \end{vmatrix}$$

$$= 9(8 - 9) + 3(-24 + 27) + 9(9 - 9)$$
$$= 9(-1) + 3(3) + 9(0)$$
$$= -9 + 9 + 0 = 0$$
and h² - ab = (-3)² - 9(1) = 9 - 9 = 0



 \therefore the given equation represents a pair of lines.

Let θ be the acute angle between the lines.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$= \left| \frac{2\sqrt{(-3)^2 - 9(1)}}{10} \right|$$
$$= \left| \frac{2\sqrt{9 - 9}}{10} \right| = 0$$

 \therefore tan θ = tan 0°

$$\therefore \Theta = 0^{\circ}.$$

Miscellaneous Exercise 4 | Q 13.2 | Page 132

Show that the following equation represents a pair of line. Find the acute angle between them:

 $2x^2 + xy - y^2 + x + 4y - 3 = 0$

Solution: Comparing this equation with

$$ax^{2} + 2hxy + by^{2} + 2gx + 2fy + c = 0$$
, we get,

a = 2, h =
$$\frac{1}{2}$$
, b = -1, g = $\frac{1}{2}$, f = 2 and c = -3
∴ D = $\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}$
= $\begin{vmatrix} 2 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -1 & 2 \\ \frac{1}{2} & 2 & -3 \end{vmatrix}$



$$= 2(3-4) - \frac{1}{2}\left(-\frac{3}{2}-1\right) + \frac{1}{2}\left(1+\frac{1}{2}\right)$$
$$= -2 + \frac{3}{4} + \frac{1}{2} + \frac{1}{2} + \frac{1}{4}$$
$$= -2 + 1 + 1$$
$$= -2 + 2 = 0$$

 \therefore the given equation represents a pair of lines.

Let θ be the acute angle between the lines.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$= \left| \frac{2\sqrt{\left(\frac{1}{2}\right)^2 - (2)(-1)}}{2 - 1} \right|$$
$$= \left| \frac{2\sqrt{\frac{1}{4} + 2}}{1} \right|$$
$$= 2\sqrt{\frac{9}{4}} = 3$$

 $\therefore \tan \theta = \tan 3$

 $\therefore \theta = \tan^{-1} (3)$

Miscellaneous Exercise 4 | Q 13.3 | Page 132

Show that the following equation represents a pair of line. Find the acute angle between them:

 $(x - 3)^2 + (x - 3)(y - 4) - 2(y - 4)^2 = 0$

Solution: Put x - 3 = X and y - 4 = Y in the given equation, we get,



$X^2 + XY - 2Y^2 = 0$

Comparing this equation with $ax^2 + 2hxy + by^2 = 0$, we get,

a = 1, h = 1/2, b = - 2

This is the homogeneous equation of second degree

and h
2
 - ab = $\left(rac{1}{2}
ight)^2 - 1(-2)$ $= rac{1}{4} + 2 = rac{9}{4} > 0$

Hence, it represents a pair of lines passing through the new origin (3, 4). Let θ be the acute angle between the lines.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$

here $a = 1, 2h = 1, i.e. h = \frac{1}{2}$ and $b = -2$
$$\therefore \tan \theta = \left| \frac{2\sqrt{\left(\frac{1}{2}\right)^2 - 1(-2)}}{1 - 2} \right|$$
$$= \left| \frac{2\left(\sqrt{\frac{1}{4} + 2}\right)}{-1} \right|$$
$$= \left| \frac{2 \times \frac{3}{2}}{-1} \right|$$
$$\therefore \tan \theta = 3$$

 $\therefore \theta = \tan^{-1}(3)$



Miscellaneous Exercise 4 | Q 14 | Page 132

Find the combined equation of lines passing through the origin and each of which making an angle of 60° with the Y-axis.

Solution:



Let OA and OB be the lines through the origin making an angle of 60° with the Y-axis. Then OA and OB make an angle of 30° and 150° with the positive direction of X-axis.

$$\therefore$$
 slope of OA = tan 30° = $\frac{1}{\sqrt{3}}$

 \therefore equation of the line OA is

$$y = \frac{1}{\sqrt{3}} x$$
 i.e. $x - \sqrt{3}y = 0$

Slope of OB = $\tan 150^\circ = \tan (180^\circ - 30^\circ)$

$$= - \tan 30^\circ = -\frac{1}{\sqrt{3}}$$

: equation of the line OB is

$$y = -\frac{1}{\sqrt{3}}x$$
 i.e. $x + \sqrt{3}y = 0$

 \therefore required combined equation is

$$\left(\mathbf{x} - \sqrt{3}\mathbf{y}\right)\left(\mathbf{x} + \sqrt{3}\mathbf{y}\right) = 0$$

i.e. $x^2 - 3y^2 = 0$


Miscellaneous Exercise 4 | Q 15 | Page 132

If the lines represented by $ax^2 + 2hxy + by^2 = 0$ make angles of equal measure with the coordinate axes, then show that $a \pm b$.

OR

Show that, one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ will make an angle of the same measure with the X-axis as the other makes with the Y-axis, if $a = \pm b$.

Solution:



Let OA and OB be the two lines through the origin represented by $ax^2 + 2hxy + by^2 = 0$.

Since these lines make angles of equal measure with the coordinate axes, they make angles α and $\pi/2$ - α with the positive direction of X-axis or α and $\pi/2 + \alpha$ with the positive direction of X-axis.

 \therefore slope of the line OA = m₁ = tan α

and slope of the line $OB = m_2$

$$= \tan\left(\frac{\pi}{2} - \alpha\right) \operatorname{or} \tan\left(\frac{\pi}{2} + \alpha\right)$$

i.e. $m_2 = \cot \alpha \text{ or } m_2 = - \cot \alpha$

$$\therefore$$
 m₁m₂ = tan α × cot α = 1



OR $m_1m_2 = \tan \alpha (-\cot \alpha) = -1$ i.e. $m_1m_2 = \pm 1$ But $m_1m_2 = ab$ $\therefore a/b=\pm 1$ $\therefore a = \pm b$ This is the required condition.

Miscellaneous Exercise 4 | Q 16 | Page 132

Show that the combined equation of the pair of lines passing through the origin and each making an angle α with the line x + y = 0 is x² + 2(sec 2\alpha)xy + y² = 0

Solution: Let OA and OB be the required lines.

Let OA (or OB) has slope m.

 \therefore its equation is y = mx ...(1)

It makes an angle α with x + y = 0 whose slope is - 1.

 $\therefore \tan \alpha = \left| \frac{m+1}{1+m(-1)} \right|$

Squaring both sides, we get,

$$an^2lpha=rac{\left(\mathrm{m}+1
ight)^2}{\left(1-\mathrm{m}
ight)^2}$$

$$\therefore \tan^2 \alpha (1 - 2m + m^2) = m^2 + 2m + 1$$

$$\therefore \tan^2 \alpha - 2 \operatorname{mtan}^2 \alpha + \operatorname{m}^2 \tan^2 \alpha = \operatorname{m}^2 + 2 \operatorname{m} + 1$$

 $\therefore (\tan^2\alpha - 1)m^2 - 2(1 + \tan^2\alpha)m + (\tan^2\alpha - 1) = 0$



$$\therefore m^{2} - 2\left(\frac{1 + \tan^{2}\alpha}{\tan^{2}\alpha - 1}\right)m + 1 = 0$$

$$\therefore m^{2} + 2\left(\frac{1 + \tan^{2}\alpha}{1 - \tan^{2}\alpha}\right)m + 1 = 0$$

$$\therefore m^{2} + 2(\sec 2\alpha)m + 1 = 0 \dots \left[\because \cos 2\alpha = \frac{1 - \tan^{2}\alpha}{1 + \tan^{2}\alpha}\right]$$

$$\therefore \frac{y^{2}}{x^{2}} + 2(\sec 2\alpha)\frac{y}{x} + 1 = 0$$

$$\therefore y^{2} 2xy \sec 2\alpha + x^{2} = 0 \dots [By (1)]$$

$$\therefore y^{2} + 2xy \sec 2\alpha + x^{2} = 0$$

$$\therefore x^{2} + 2(\sec 2\alpha)xy + y^{2} = 0 \text{ is the required equation.}$$

Miscellaneous Exercise 4 | Q 17 | Page 132

Show that the line 3x + 4y + 5 = 0 and the lines $(3x + 4y)^2 - 3(4x - 3y)^2 = 0$ form the sides of an equilateral triangle.

Solution: The slope of the line 3x + 4y + 5 = 0 is $m_1 = -3/4$

Let m be the slope of one of the line making an angle of 60° with the line 3x + 4y + 5 = 0. The angle between the lines having slope m and m₁ is 60° .

$$\therefore \tan 60^\circ = \left| \frac{\mathbf{m} - \mathbf{m}_1}{1 + \mathbf{m} \cdot \mathbf{m}_1} \right|, \text{ where } \tan 60^\circ = \sqrt{3}$$
$$\therefore \sqrt{3} = \left| \frac{\mathbf{m} - \left(-\frac{3}{4}\right)}{1 + \mathbf{m}\left(-\frac{3}{4}\right)} \right|$$
$$\therefore \sqrt{3} = \left| \frac{4\mathbf{m} + 3}{4 - 3\mathbf{m}} \right|$$



On squaring both sides, we get,

$$3 = \frac{(4m + 3)^2}{(4 - 3m)^2}$$

$$\therefore 3(4 - 3m)^2 = (4m + 3)^2$$

$$\therefore 3(16 - 24m + 9m^2) = 16m^2 + 24m + 9$$

$$\therefore 48 - 72m + 27m^2 = 16m^2 + 24m + 9$$

$$\therefore 11m^2 - 96m + 39 = 0$$

This is the auxiliary equation of the two lines and their joint equation is obtained by putting m = y/x.

 \therefore the combined equation of the two lines is

$$11\left(\frac{y}{x}\right)^2 - 96(y'/x) + 39 = 0$$
$$\therefore \frac{11y^2}{x^2} - \frac{96y}{x} + 39 = 0$$

- $\therefore 11y^2 96xy + 39x^2 = 0$
- $\therefore 39x^2 96xy + 11y^2 = 0$

 \therefore 39x² - 96xy + 11y² = 0 is the joint equation of the two lines through the origin each making an angle of 60° with the line 3x + 4y + 5 = 0

The equation $39x^2 - 96xy + 11y^2 = 0$ can be written as: $-39x^2 + 96xy - 11y^2 = 0$

i.e.
$$(9x^2 - 48x^2) + (24xy + 72xy) + (16y^2 - 27y^2) = 0$$

i.e. $(9x^2 + 24xy + 16y^2) - 3(16x^2 - 24xy + 9y^2) = 0$

i.e. $(3x + 4y)^2 - 3(4x - 3y)^2 = 0$

Hence, the line 3x + 4y + 5 = 0 and the lines $(3x + 4y)^2 - 3(4x - 3y)^2$ form the sides of an equilateral triangle.

Miscellaneous Exercise 4 | Q 18 | Page 132

Show that the lines $x^2 - 4xy + y^2 = 0$ and the line $x + y = \sqrt{6}$ form an equilateral triangle.

Find its area and perimeter.

Solution: $x^2 - 4xy + y^2 = 0$ and $x + y = \sqrt{6}$ form a triangle OAB which is equilateral.



Let OM be the perpendicular from the origin O to AB whose equation is $x + y = \sqrt{6}$



$$\sin 60^\circ = \frac{OM}{OA}$$
$$\therefore \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{OA}$$

 \therefore length of the each side of the equilateral triangle OAB = 2 units.

 \therefore perimeter of \triangle OAB = 3 × length of each side

 $= 3 \times 2 = 6$ units

Miscellaneous Exercise 4 | Q 19 | Page 132

If the slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is square of the slope of the other line, show that $a^2b + ab^2 + 8h^3 = 6abh$.



Solution: Let m be the slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$.

Then the other line has slope m²

$$\therefore m + m^{2} = \frac{-2h}{b} \qquad \dots (1) \text{ and}$$

$$(m)(m^{2}) = \frac{a}{b}$$
i.e. $m^{3} = \frac{a}{b} \qquad \dots (2)$

$$\therefore (m + m^{2})^{3} = m^{3} + (m^{2})^{3} + 3(m)(m^{2})(m + m^{2}) \dots [\because (p + q)^{3} = p^{3} + q^{3} + 3pq(p + q)]$$

$$\therefore \left(\frac{-2h}{b}\right)^{3} = \frac{a}{b} + \frac{a^{2}}{b^{2}} + 3\frac{a}{b}\left(\frac{-2h}{b}\right)$$

$$\therefore \frac{-8h^{3}}{b^{3}} = \frac{a}{b} + \frac{a^{2}}{b^{2}} - \frac{6ah}{b^{2}}$$

Multiplying by b³, we get,

 $-8h^3 = ab^2 + a^2b - 6abh$

 $\therefore a^2b + ab^2 + 8h^3 = 6abh$

This is the required condition.

Miscellaneous Exercise 4 | Q 20 | Page 132

Prove that the product of length of perpendiculars drawn from

$$\begin{aligned} \mathsf{P}(\mathsf{x}_1, \mathsf{y}_1) \text{ to the lines represented by } \mathsf{a}\mathsf{x}^2 + 2\mathsf{h}\mathsf{x}\mathsf{y} + \mathsf{b}\mathsf{y}^2 &= 0 \text{ is} \\ \left| \frac{\mathsf{a}\mathsf{x}_1^2 + 2\mathsf{h}\mathsf{x}_1\mathsf{y}_1 + \mathsf{b}\mathsf{y}_1^2}{\sqrt{\mathsf{a}-\mathsf{b}}^2 + 4\mathsf{h}^2} \right| \end{aligned}$$

Solution: Let m_1 and m_2 be the slopes of the lines represented by $ax^2 + 2hxy + by^2 = 0$



$$\therefore$$
 m₁ + m₂ = $-\frac{2\mathbf{h}}{\mathbf{b}}$ and m₁m₂ = $\frac{\mathbf{a}}{\mathbf{b}}$...(1)

The separate equations of the lines represented by $ax^2 + 2hxy + by^2 = 0$ are

$$y = m_1 x$$
 and $y = m_2 x$

i.e.
$$m_1x - y = 0$$
 and $m_2x - y = 0$

Length of perpendicular from $P(x_1, y_1)$ on

$$m_1 x - y = 0$$
 is $\left| \frac{m_1 x_1 - y_1}{\sqrt{m_1^2 + 1}} \right|$

Length of perpendicular form $P(x_1, y_1)$ on

$$m_2 x - y = 0$$
 is $\left| \frac{m_2 x_1 - y_1}{\sqrt{m_2^2 + 1}} \right|$

... product of lengths of perpendiculars

$$egin{aligned} &= \left|rac{\mathrm{m}_1 \mathrm{x}_1 - \mathrm{y}_1}{\sqrt{\mathrm{m}_1^2 + 1}}
ight| imes \left|rac{\mathrm{m}_2 \mathrm{x}_1 - \mathrm{y}_1}{\sqrt{\mathrm{m}_2^2 + 1}}
ight| \ &= \left|rac{\mathrm{m}_1 \mathrm{m}_2 \mathrm{x}_1^2 - (\mathrm{m}_1 + \mathrm{m}_2) \mathrm{x}_1 \mathrm{y}_1 + \mathrm{y}_1^2}{\sqrt{\mathrm{m}_1^2 \mathrm{m}_2^2 + \mathrm{m}_1^2 + \mathrm{m}_2^2 + 1}}
ight| \end{aligned}$$



$$=rac{\mathrm{m_1m_2x_1^2}-(\mathrm{m_1+m_2})\mathrm{x_1y_1}+\mathrm{y_1^2}}{\sqrt{\mathrm{m_1^2m_2^2}+(\mathrm{m_1+m_2})^2-2\mathrm{m_1m_2}+1}}$$

$$= \left| \frac{\frac{a}{b} \cdot x_1^2 - \frac{-2h}{b} x_1 y_1 + y_1^2}{\sqrt{\frac{a^2}{b^2} + \frac{-2h}{b} - \frac{2a}{b} + 1}} \right| \quad ...(\text{By (1)})$$

$$= \left| rac{{{\mathrm{ax}}_1^2 + 2{{\mathrm{hx}}_1}{\mathrm{y}_1} + {\mathrm{by}}_1^2 }}{{\sqrt {{\mathrm{a}}^2 + 4{{\mathrm{h}}^2} - 2{\mathrm{ab}} + {\mathrm{b}}^2 }}}
ight|$$

$$egin{aligned} &= \left| rac{\mathrm{ax}_1^2 + 2\mathrm{hx}_1\mathrm{y}_1 + \mathrm{by}_1^2}{\sqrt{\left(\mathrm{a}^2 - 2\mathrm{ab} + \mathrm{b}^2
ight) + 4\mathrm{h}^2}}
ight| \ &= \left| rac{\mathrm{ax}_1^2 + 2\mathrm{hx}_1\mathrm{y}_1 + \mathrm{by}_1^2}{\sqrt{\left(\mathrm{a} - \mathrm{b}
ight)^2 + 4\mathrm{h}^2}}
ight| \end{aligned}$$

Miscellaneous Exercise 4 | Q 21 | Page 132

Show that the difference between the slopes of the lines given by $(\tan^2\theta + \cos^2\theta)x^2 - 2xy \tan \theta + (\sin^2\theta)y^2 = 0$ is two.

Solution: Comparing the equation

 $(\tan^2\theta + \cos^2\theta)x^2 - 2xy \tan \theta + (\sin^2\theta)y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, wew get,

a = $tan^2\theta + cos^2\theta$, 2h = - 2tan θ b = $sin^2\theta$

Let m_1 and m_2 be the slopes of the lines represented by the given equation.



$$\therefore m_1 + m_2 = \frac{-2h}{b} = -\left[\frac{-2\tan\theta}{\sin^2\theta}\right] = \frac{2\tan\theta}{\sin^2\theta} \quad \dots (1)$$

and $m_1m_2 = \frac{a}{b} = \frac{\tan^2\theta + \cos^2\theta}{\sin^2}\theta \quad \dots (2)$
$$\therefore (m_1 - m_2)^2 = (m_1 + m_2)^2 - 4m_1m_2$$

$$= \left(\frac{2\mathrm{tan}\theta}{\mathrm{sin}^2\theta}\right)^2 - 4\left(\frac{\mathrm{tan}^2\theta + \mathrm{cos}^2\theta}{\mathrm{sin}^2\theta}\right)$$

$$=rac{4 an^2 heta}{\sin^4 heta}-4igg(rac{ an^2 heta+\cos^2 heta}{\sin^2 heta}igg)\ =rac{4igg(rac{\sin^2 heta}{\cos^2 heta}igg)}{\sin^4 heta}-4igg[rac{igg(rac{\sin^2 heta+\cos^2 heta}{\cos^2 heta}+\cos^2 hetaigg)}{\sin^2 heta}igg]$$

$$=\frac{4}{\sin^2\theta.\cos^2\theta}-\frac{4\left(\sin^2\theta+\cos^4\theta\right)}{\sin^2\theta.\cos^2\theta}$$

$$=4\left[\frac{1-\sin^2\theta-\cos^4\theta}{\sin^2\theta.\cos^2\theta}\right]$$

$$=4\left[\frac{\cos^2\theta-\cos^4\theta}{\sin^2\theta.\cos^2\theta}\right]$$



$$=4\left[rac{\cos^2 hetaig(1-\cos^2 hetaig)}{\sin^2 heta.\cos^2 heta}
ight]=4$$

 $|m_1 - m_2| = 2$

 \therefore the slopes differ by 2.

Miscellaneous Exercise 4 | Q 22 | Page 132

Find the condition that the equation $ay^2 + bxy + ex + dy = 0$ may represent a pair of lines.

Solution: Comparing the equation $ay^2 + bxy + ex + dy = 0$ with $Ax^2 + 2Hxy + By^2 + 2Gx + 2Fy + C = 0$, we get,

$$A = 0, H = rac{b}{2}, B = a, G = rac{e}{2}, F = rac{d}{2}, C = 0$$

The given equation represents a pair of lines,

$$\begin{vmatrix} A & H & G \\ H & B & F \\ G & F & C \end{vmatrix} = 0$$

i.e. if
$$\begin{vmatrix} 0 & \frac{b}{2} & \frac{e}{2} \\ \frac{b}{2} & a & \frac{d}{2} \\ \frac{e}{2} & \frac{d}{2} & 0 \end{vmatrix} = 0$$

i.e. if $0 - \frac{b}{2} \left(0 - \frac{ed}{4} \right) + \frac{e}{2} \left(\frac{bd}{4} - \frac{ae}{2} \right) = 0$
i.e. if $\frac{bed}{8} + \frac{bed}{8} - \frac{ae^2}{4} = 0$
i.e. if bed - ae² = 0
i.e. if e(bd - ae) = 0
i.e. if e = 0 or bd - ae = 0
i.e. if e = 0 or bd = ae



This is the required condition.

Miscellaneous Exercise 4 | Q 23 | Page 132

If the lines given by $ax^2 + 2hxy + by^2 = 0$ form an equilateral triangle with the line lx + my = 1, show that $(3a + b)(a + 3b) = 4h^2$.

Solution: Since the lines $ax^2 + 2hxy + by^2 = 0$ form an equilateral triangle with the line lx + my = 1, the angle between the lines

$$ax^{2} + 2hxy + by^{2} = 0$$
 is 60°.

$$\therefore \tan 60^\circ = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
$$\therefore \sqrt{3} = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$

$$\therefore 3(a + b)^{2} = 4(h^{2} - ab)$$

$$\therefore 3(a^{2} + 2ab + b^{2}) = 4h^{2} - 4ab$$

$$\therefore 3a^{2} + 6ab + 3b^{2} + 4ab = 4h^{2}$$

$$\therefore 3a^{2} + 10ab + 3b^{2} = 4h^{2}$$

$$\therefore 3a^{2} + 9ab + ab + 3b^{2} = 4h^{2}$$

$$\therefore 3a(a + 3b) + b(a + 3b) = 4h^{2}$$

$$\therefore (3a + b)(a + 3b) = 4h^{2}$$

This is the required condition.

Miscellaneous Exercise 4 | Q 24 | Page 132

If the line x + 2 = 0 coincides with one of the lines represented by the equation $x^2 + 2xy + 4y + k = 0$, then prove that k = -4. **Solution:** One of the lines represented by $x^2 + 2xy + 4y + k = 0$ is x + 2 = 0.(1) Let the other line represented by (1) be ax + by + c = 0 \therefore their combined equation is (x + 2)(ax + by + c) = 0 $\therefore ax^2 + bxy + cx + 2ax + 2by + 2c = 0$

$$\therefore ax^2 + bxy + (2a + c)x + 2by + 2c = 0$$
 ...(2)



As the equations (1) and (2) are the combined equations of the same two lines, they are identical.

.. by comparing their corresponding coefficients, we get,

$$\frac{a}{1} = \frac{b}{2} = \frac{2b}{4} = \frac{2c}{k} \text{ and } 2a + c = 0$$

$$\therefore a = \frac{2c}{k} \text{ and } c = -2a$$

$$\therefore a = \frac{2(-2a)}{k}$$

$$\therefore 1 = \frac{-4}{k}$$

$$\therefore k = -4$$

Miscellaneous Exercise 4 | Q 25 | Page 132

Prove that the combined of the pair of lines passing through the origin and perpendicular to the lines $ax^2 + 2hxy + by^2 = 0$ is $bx^2 - 2hxy + ay^2 = 0$. **Solution:** Let m₁ and m₂ be the slopes of the lines represented by $ax^2 + 2hxy + by^2 = 0$.

$$\therefore$$
 m₁ + m₂ = $\frac{-2\mathbf{h}}{\mathbf{b}}$ and m₁m₂ = $\frac{\mathbf{a}}{\mathbf{b}}$ (1)

Now, the required lines are perpendicular to these lines.

$$\therefore$$
 their slopes are $-\frac{1}{m_1}$ and $-\frac{1}{m_2}$

Since these lines are passing through the origin, their separate equations are

$$\mathsf{y} = -\frac{1}{m_1}\mathsf{x} \quad \text{and} \ \mathsf{y} = -\frac{1}{m_2}\mathsf{x}$$

i.e. $m_1y = -x$ and $m_2y = -x$

i.e. $x + m_1 y = 0$ and $x + m_2 y = 0$

 \therefore their combined equation is

 $(x + m_1 y)(x + m_2 y) = 0$

$$\therefore x^2 + (m_1 + m_2)xy + m_1m_2y^2 = 0$$



$$\therefore x^2 \frac{-2h}{b} x + \frac{a}{b} y^2 = 0 \quad ...[By(1)]$$

 $\therefore bx^2 - 2hxy + ay^2 = 0$

Miscellaneous Exercise 4 | Q 26 | Page 132

If equation $ax^2 - y^2 + 2y + c = 1$ represents a pair of perpendicular lines, then find a and c.

Solution: The given equation represents a pair of lines perpendicular to each other.

: coefficient of
$$x^2$$
 + coefficient of $y^2 = 0$

With this value of a, the given equation is

 $x^2 - y^2 + 2y + c - 1 = 0$

Comparing this equation with

 $Ax^{2} + 2Hxy + By^{2} + 2Gx + 2Fy + C = 0$, we get,

A = 1, H = 0, B = -1, G = 0, F = 1, C = c - 1

Since the given equation represents a pair of lines,

$$D = \begin{vmatrix} A & H & G \\ H & B & F \\ G & F & C \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} 1 & 0 & 0 \\ 0 & -1 & 1 \\ 0 & 1 & c - 1 \end{vmatrix} = 0$$

$$\therefore 1(-c + 1 - 1) - 0 + 0 = 0$$

$$\therefore -c = 0$$

Hence, $a = 1, c = 0$.